



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

T cell–intrinsic prostaglandin E₂-EP2/EP4 signaling is critical in pathogenic TH17 cell–driven inflammation

Citation for published version:

Lee, J, Aoki, T, Thumkeo, D, Siriwach, R, Yao, C & Narumiya, S 2019, 'T cell–intrinsic prostaglandin E₂-EP2/EP4 signaling is critical in pathogenic T_H17 cell–driven inflammation', *Journal of Allergy and Clinical Immunology*, vol. 143, no. 2, pp. 631-643. <https://doi.org/10.1016/j.jaci.2018.05.036>

Digital Object Identifier (DOI):

[10.1016/j.jaci.2018.05.036](https://doi.org/10.1016/j.jaci.2018.05.036)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Allergy and Clinical Immunology

Publisher Rights Statement:

Open Access funded by Medical Research Council
Under a Creative Commons license

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Accepted Manuscript

T cell-intrinsic prostaglandin E₂-EP2/EP4 signaling is critical in pathogenic Th17 cell-driven inflammation

Jinju Lee, MSc, Tomohiro Aoki, MD, PhD, Dean Thumkeo, MD, PhD, Ratklao Siriwach, PhD, Chengcan Yao, PhD, Shuh Narumiya, MD, PhD

PII: S0091-6749(18)30896-0

DOI: [10.1016/j.jaci.2018.05.036](https://doi.org/10.1016/j.jaci.2018.05.036)

Reference: YMAI 13487

To appear in: *Journal of Allergy and Clinical Immunology*


Received Date: 18 December 2017

Revised Date: 3 May 2018

Accepted Date: 25 May 2018


Please cite this article as: Lee J, Aoki T, Thumkeo D, Siriwach R, Yao C, Narumiya S, T cell-intrinsic prostaglandin E₂-EP2/EP4 signaling is critical in pathogenic Th17 cell-driven inflammation, *Journal of Allergy and Clinical Immunology* (2018), doi: 10.1016/j.jaci.2018.05.036.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.





Intrinsic PGE₂ drives Th17 pathogenicity.



PGE₂, Prostaglandin E₂; COX2, cyclooxygenase 2; cAMP, cyclic adenosine monophosphate; PKA, Protein kinase A; CREB, cAMP response element binding protein; Jak2, Janus kinase 2; STAT3, Signal transducer and activator of transcription 3.

**T cell-intrinsic prostaglandin E₂-EP2/EP4 signaling is critical
in pathogenic Th17 cell-driven inflammation**

Jinju Lee, MSc^{a,b}, Tomohiro Aoki, MD, PhD^{a,c}, Dean Thumkeo, MD, PhD^{a,c}, Ratklao
Siriwach, PhD^c, Chengcan Yao, PhD^d, Shuh Narumiya, MD, PhD^{a,c}

^a Core Research for Evolutional Science and Technology (CREST), Medical Innovation
Center, Kyoto University Graduate School of Medicine, Kyoto 606-8507, Japan.

^b Kyoto University, Graduate School of Biostudies, Kyoto 606-8501, Japan

^c Center for Innovation in Immunoregulation Technology and Therapeutics, Kyoto
University Graduate School of Medicine, Kyoto 606-8507, Japan.

^d Medical Research Council (MRC) Centre for Inflammation Research, Queen's
Medical Research Institute, The University of Edinburgh, Edinburgh EH16 4TJ, UK.

Corresponding authors:

Shuh Narumiya, MD, PhD (ORCID ID: 0000-0001-8062-6529)

Medical Innovation Center, Kyoto University Graduate School of Medicine

53 Shogoin Kawaharacho, Sakyo, Kyoto 606-8507, Japan

Telephone: +81-75-366-7468, Fax: +81-75-753-9114

E-mail: snaru@mfour.med.kyoto-u.ac.jp

Chengcan Yao, PhD (ORCID ID: 0000-0003-3754-2842)

Medical Research Council (MRC) Centre for Inflammation Research,

Queen's Medical Research Institute, The University of Edinburgh

47 Little France Crescent, Edinburgh EH16 4TJ, UK.

Telephone: +4413-1242-6685, Fax: +4413-1242-6578

E-mail: Chengcan.Yao@ed.ac.uk

Funding:

This work was supported in part by Core Research for Evolutional Science and
Technology Program on Chronic Inflammation (15gm0410006h0006) from the Japan
Agency for Medical Research and Development (S.N.), by a Coordination Fund from
the Japanese Ministry for Education, Culture, Sports, Science and Technology (MEXT)
and Astellas Pharma Inc. to Kyoto University (S.N.), by a collaborative grant to the
Kyoto University from Ono Pharmaceuticals (S.N.), and by Medical Research Council
UK (MR/R008167/1 to C.Y.), Cancer Research UK (C63480/A25246 to C.Y.) and
Wellcome Trust Institutional Strategic Support Fund (C.Y.).

Disclosure of potential conflict of interest:

T.A., D.T., and S.N. were supported by the Coordination Fund from MEXT and Astellas
Pharma Inc. S.N. is a scientific advisor to Astellas Pharma Inc. Other authors declare
that they have no conflict of interests.

ABSTRACT

Background: Interleukin-23 (IL-23) is the key cytokine for generation of pathogenic IL-17-producing helper T (Th17) cells that critically contribute to autoimmune diseases. However, how IL-23 generates pathogenic Th17 cells remains to be elucidated.

Objectives: To examine the involvement, molecular mechanisms and clinical implications of prostaglandin (PG) E₂-EP2/EP4 signaling in induction of IL-23-driven pathogenic Th17 cells.

Methods: The role of PGE₂ in induction of pathogenic Th17 cells was investigated in mouse Th17 cells in culture *in vitro* and in IL-23-induced psoriasis mouse model *in vivo*. Clinical relevance of findings in mice was examined by gene expression profiling of IL-23 and PGE₂-EP2/EP4 signaling in psoriatic skin from patients.

Results: IL-23 induces *ptgs2* encoding cyclooxygenase 2 in Th17 cells and produces PGE₂, which acts back on PGE receptors EP2 and EP4 in these cells and enhances IL-23-induced expression of an IL-23 receptor subunit gene, *Il23r*, by activating STAT3, CREB1 and NF-κB through cAMP-protein kinase A signaling. This PGE₂ signaling also induces expression of various inflammation-related genes, which possibly function in Th17 cell-mediated pathology. Combined deletion of EP2 and EP4 selectively in T cells suppressed accumulation of IL-17A⁺ and IL-17A⁺IFN-γ⁺ pathogenic Th17 cells and abolished skin inflammation in IL-23-induced psoriasis mouse model. Analysis of human psoriatic skin biopsies shows positive correlation between PGE₂ signaling and the IL-23/Th17 pathway.

Conclusions: The T cell-intrinsic EP2/EP4 signaling is critical in IL-23-driven generation of pathogenic Th17 cells and consequent pathogenesis in the skin.

Key Messages

IL-23 triggers T cell-intrinsic PGE₂-EP2/EP4 signaling that is critical in Th17 cell-mediated immune pathogenesis.

The PGE₂-EP2/EP4 signaling functions synergistically with IL-23 and not only amplifies *Il23r* expression but also induces a unique pathogenic gene expression signature by activating STAT3, CREB1 and NF-κB.

This PGE₂ signaling can be a therapeutic target of Th17 cell-mediated diseases, because combined blockade of EP2 and EP4 suppresses IL-23-induced pathogenic Th17 cell generation and consequent psoriatic skin inflammation.

Capsule Summary

IL-23 mobilizes T cell-intrinsic PGE₂-EP2/EP4 signaling, which is critical in IL-23-induced pathogenic Th17 cell generation. Combined blockade of EP2 and EP4 suppressed IL-23-induced skin inflammation, suggesting this pathway as potential therapeutic target of Th17-mediated diseases.

Key words: *psoriasis, pathogenic Th17 cells, IL-23R, prostaglandin E₂, prostaglandin E receptor EP2, prostaglandin E receptor EP4, STAT3, CREB1, NF-κB*

Abbreviations used:

Th1, helper T cell type 1

Th2, helper T cell type 2

Th17, helper T cell type 17

Treg, regulatory T cell

PG, prostaglandin

- 90 COX, cyclooxygenase
- 91 cAMP, cyclic adenosine monophosphate
- 92 PKA, protein kinase A
- 93 CREB, cAMP Responsive Element Binding Protein
- 94 Epac, exchange factor directly activated by cAMP
- 95 db-cAMP, dibutyryl cAMP
- 96 IMQ, imiquimod

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

INTRODUCTION

CD4⁺ T cells differentiate into Th1, Th2 and Th17 cells in response to specific cytokine milieu present in microenvironment of inflammation and mediate immune inflammatory responses in respective settings.¹⁻⁴ Among these Th subsets, Th17 cells mediate inflammatory responses in many autoimmune diseases including multiple sclerosis, inflammatory bowel diseases such as Crohn's disease, psoriasis and rheumatoid arthritis. The importance of Th17 cells in these processes was suggested first in animal models of these diseases including experimental autoimmune encephalomyelitis (EAE) and IL-23- or imiquimod-induced psoriasis model,⁵⁻⁹ and validated recently by clinical effectiveness of antibodies targeting to IL-23 in patients with psoriasis.¹⁰⁻¹⁴

Differentiation of Th17 cells from naïve CD4⁺ T cells is driven by the combined actions of interleukin-6 (IL-6) and transforming growth factor- β 1 (TGF- β 1).¹⁵⁻¹⁹ However, differentiated Th17 cells have little capacity to induce autoimmune and inflammatory pathology.²⁰ It should be noted that these Th17 cells exhibit plasticity and could transdifferentiate into other effector T cell types or even regulatory T cells under certain context such as inflammation or autoimmune disease.²¹⁻²³ Accumulating evidences suggest that T cell intrinsic IL-23 signaling not only increases IL-17 production of Th17 cells but also plays a crucial role in inducing and stabilizing their pathogenicity.^{20,24-27} It is known that IL-23 acts on IL-23 receptor composed of IL-23R and IL-12R β 1, activates signal transducer and activator of transcription 3 (STAT3) and induces expression of *Il23r*, thus forming the self-amplification loop. The pathophysiological importance of this IL-23-IL-23 receptor signaling has been indicated by several genomic studies that showed positive correlation between single nucleotide

polymorphisms (SNPs) of genes involved in this pathway, e.g., *IL23R*, *IL12B* (*p40*), *JAK2* and *STAT3*, and a wide range of IL-17-dependent autoimmune diseases.²⁸⁻³⁰

While it was shown that IL-23 signaling induces expression of Th17 pathogenic signature genes through activation of *STAT3*,^{31,32} other transcription factor(s) besides *STAT3* are also implicated for induction of pathogenic Th17 cells, because IL-6 that activates *STAT3* similarly to IL-23 cannot induce *IL-23R* gene expression.³² The identity of additional transcriptional factor(s) and regulatory mechanisms are therefore important issues to be defined. Moreover, how IL-23 cooperates with other inflammatory factors formed in disease microenvironment and how critical is such cooperation for pathogenic conversion of Th17 cells and overall pathology remain largely obscure. Clarification of these points could provide a new opportunity to develop small molecule drugs as therapeutic alternatives to anti-IL-23 antibodies without systemic immune suppression. Biological agents may additionally cause unpredictable adverse events³³ and can be costly on long-term use.³⁴ It should also be mentioned that JAK inhibitors that are now being evaluated in their efficacy in autoimmune diseases are presumably not free from adverse effects, either, because of their effects of general immune suppression.³⁵

Prostanoids including prostaglandin (PG) D_2 , PGE_2 , $PGF_{2\alpha}$, PGI_2 and thromboxane A_2 (TXA_2) are oxygenated metabolites of arachidonic acid produced by sequential actions of cyclooxygenase (COX) and respective synthases, and act on their cognate receptors, DP for PGD_2 , EP1 to 4 for PGE_2 , IP for PGI_2 , FP for $PGF_{2\alpha}$, and TP for TXA_2 , to exert their actions.³⁶ While prostanoids were previously regarded as immunosuppressants,^{37,38} recent studies have revealed their immunostimulatory actions in processes such as cytokine production, dendritic cell maturation, macrophage

activation, and differentiation and expansion of Th subsets.³⁹⁻⁴¹ Indeed, the PGE₂-EP2 and EP4 (EP2/EP4) signaling enhances Th1 differentiation by inducing the expression of an IL-12 receptor subunit, *Il12rb2*, and interferon- γ receptor, *Ifngr1*, thus facilitating IL-12 signaling and Th1 differentiation.^{42,43} Notably, this PGE₂-EP2/EP4 signaling was also reported to synergize with IL-23 to facilitate Th17 cell expansion both in murine and human T cells.⁴³⁻⁴⁵ However, whether the PGE₂-EP2/EP4 signaling is involved in induction of pathogenic Th17 cells, and, if so, how remains unknown.

In this study, we have examined how the PGE₂-EP2/EP4 signaling and IL-23 stimulation together regulate the generation of pathogenic Th17 cells. Through this analysis, we have identified the transcription mechanisms in addition to STAT3 that regulate *Il23r* expression and Th17 pathogenicity. We have further clarified the importance of the PGE₂ signaling in the Th17-mediated immune inflammation *in vivo*, and found the correlation between PGE₂-EP2/EP4 signaling and IL-23-IL-23 receptor signaling in biopsy samples from psoriasis patients.

METHODS

Mice

All animal experiments were approved by the Institutional Animal Care and Use Committee of Kyoto University Graduate School of Medicine, and complied with the National Institutes of Health Guide for the Care and Use of Laboratory Animals. C57BL/6NCrSlc mice were purchased from Shimizu laboratory, and Lck-Cre mice and B6.Cg-*Nfkb1*^{tm1Bal}/J mice were purchased from Jackson Laboratory. Mice deficient in *Ptger2*⁴⁶ and mice with floxed *Ptger2*⁴⁷ were established in our laboratory. Mice with floxed *Ptger4* was a kind gift of Richard Breyer.⁴⁸

Psoriasis models

To induce psoriasis-like lesion in the ear in IL-23-induced psoriasis mouse model, mice were subcutaneously injected with IL-23 (500 ng; #130-096-677, Miltenyi, Bergisch Gladbach, Germany) once a day in one ear and with PBS in the contralateral ear as a control. In imiquimod (IMQ)-induced psoriasis mouse model, baselna cream containing 10% IMQ was applied onto the ear of mice once a day. Ear thickness was then measured by a digital micrometer (#KM-BMB1-25, Mitutoyo, Kawasaki, Japan) every other day. In some experiments, an antagonist for EP4, AS1954813,⁴⁹ suspended in 0.5 % methylcellulose was orally administered twice a day or indomethacin and SC-236 were administered by drinking water during the experimental period.

Other methods

See the Supplementary Methods section in this article's Online Repository at www.jacionline.org.

RESULTS

IL-23 mobilizes the endogenous COX-2-PGE₂-EP2/EP4 signaling that enhances induction of *Il23r* expression in Th17 cells

Given the previous findings⁴³⁻⁴⁵ that the PGE₂-EP2/EP4 signaling enhances IL-23-induced Th17 cell expansion, we questioned here whether and how this signaling contributes to pathogenic Th17 cell generation by IL-23. To investigate this issue, we first cultured CD4⁺ T cells from mouse spleen under the Th17-skewing condition (IL-6 plus TGF- β 1) for 4 days, and then incubated with IL-23 for additional 3 days. Consistent with our previous findings,⁴³ the addition of PGE₂ to the latter culture significantly enhanced IL-23-induced expansion and *Il17a* expression of Th17 cells (Fig. 1, A and B). Interestingly, we also noted that PGE₂ markedly up-regulated IL-23-induced expression of *Il23r*, which was mimicked by both EP2- and EP4-selective agonists (Fig. 1, C). Since both EP2 and EP4 activates PKA and Epac by increasing intracellular cAMP,³⁶ we examined effects of compounds acting on these signaling, and found that a cAMP analogue, dibutyryl cAMP (db-cAMP), forskolin (FSK) and a phosphodiesterase inhibitor, IBMX, all synergized with IL-23 and significantly amplified IL-23-induced *Il23r* expression and IL-17A production in these cells (Fig. 1, D and E). Furthermore, the enhancement of *Il23r* expression was reproduced by a PKA agonist (N6-Bnz-cAMP, 300 μ M) but not an Epac activator (8-pCTP-2'-O-Me-cAMP, 300 μ M) (Fig. 1, F), and, consistently, was ameliorated by treatment with a PKA inhibitor (H-89, 10 μ M) (Fig. 1, G).

Notably, IL-23 stimulation significantly increased *Ptgs2* (COX-2) gene expression in Th17 cells (Fig. 2, A) and produced subnanomolar concentrations of PGE₂

in the culture medium (Fig. 2, B). Moreover, incubation with non-selective COX inhibitor (indomethacin, 100 μ M) or a selective COX-2 inhibitor (SC-236, 100 μ M) but not a selective COX-1 inhibitor (SC-560, 100 μ M) significantly blocked the induction of *Il23r* expression in response to both IL-23 alone and IL-23 and PGE₂ in combination (Fig. 2, C and Fig E1, A). In addition, antagonists selective to EP2 (PF-04418948) or EP4 (ONO-AE3-208) also suppressed *Il23r* expression (Fig. 2, D). Intriguingly indomethacin and SC-236 suppressed the expression of *Il23r* induced by IL-23 and PGE₂ to the level that these inhibitors achieved in the presence of IL-23 alone, suggesting that they cancelled the effect of exogenously added PGE₂ (Fig. 2, D and Fig E1, A). Given that PGE₂ added to the culture medium time-dependently degrades,⁵⁰ these results suggest that exogenously added PGE₂ induces COX-2 and produces PGE₂ endogenously and continuously as we reported previously,⁵¹ which makes more contribution to *Il23r* induction, and that indomethacin and COX-2 inhibitor block this process. Indeed, the addition of stable EP2 and EP4 agonists overcame the *Il23r* suppression by indomethacin (Fig E1, B). Therefore, these data together suggest that IL-23 stimulates Th17 cells to produce PGE₂, which acts back to EP2 and EP4 on these cells to augment *Il23r* expression in a positive feedback manner.

Induction of *Il23r* expression by IL-23 and PGE₂-cAMP signaling is mediated through not only STAT3 but also CREB1 and NF- κ B

We then investigated transcription factors responsible for induction of *Il23r* expression in Th17 cells by IL-23 and PGE₂-EP2/EP4 signaling. Because IL-23 activates STAT3 to induce *Il23r* expression,⁵² we first examined the effect of a STAT3 inhibitor. The

addition of STAT3 inhibitor VII suppressed *Il23r* expression not only by IL-23, but also by db-cAMP, and both (Fig. 3, A), indicating that the db-cAMP action was also mediated by STAT3. Consistently, Y705 phosphorylation of STAT3 was increased by db-cAMP at 5 and 30 min (Fig E2, A), which were ameliorated not only by the addition of STAT3 inhibitor VII but also by H-89 (Fig. 3, B), indicating the involvement of PKA in db-cAMP-mediated Y705 phosphorylation of STAT3. Intriguingly, the Y1007/1008 phosphorylation of JAK2, a kinase responsible for STAT3 Y705 phosphorylation in Th17 cells, was enhanced by db-cAMP, and this enhancement was suppressed by Src Kinase Inhibitor I (Fig E2, B), indicating cAMP-PKA activates STAT3 through c-Src-JAK2 pathway.

Although the above findings demonstrated that IL-23 and PGE₂-cAMP signaling converge at STAT3 activation, it is well known that other STAT3 activators, such as IL-6 and IL-21, cannot substitute for IL-23 in the expansion of Th17 population,³² indicating that STAT3 is not the sole transcription factor regulating expression of *Il23r*. Since PKA activates CREB1,³⁶ we investigated the involvement of CREB1 in *Il23r* expression. Both KG-501, a CREB1 inhibitor,⁵³ and RNAi for CREB1 suppressed *Il23r* induction in response to db-cAMP, IL-23 or both, suggesting the involvement of CREB1 in *Il23r* expression in Th17 cells (Fig. 3, C and D). As IL-23 signaling enhances endogenous PGE₂ production via induction of COX-2 expression in Th17 cells (Fig. 2, A and B), the suppression of *Il23r* expression by inhibition or depletion of CREB1 could be due to inhibition of this endogenous PGE₂ signaling for *Il23r* induction.

Furthermore, we detected an increase in S536 phosphorylation of NF-κB p65 (pp65) in response to db-cAMP, IL-23 or both at 24 h (Fig. 3, E) and an increase in

S933 phosphorylation of NF- κ B p105 subunit, a precursor of p50, in response to db-cAMP alone and its combination with IL-23 in Th17 cells (Fig. 3, E). The latter is consistent with our previous finding in dendritic cells that PGE₂-cAMP signaling activates the p50 subunit⁵⁴ and a report that phosphorylation of p105 S933 is PKA-dependent.⁵⁵ We therefore examined the involvement of NF- κ B in *Il23r* induction by using *Nfkb1*-deficient mice (p105 KO) or CTP-NBD, a NF- κ B inhibitor. Interestingly, both genetic deficiency and pharmacological inhibition of NF- κ B suppressed *Il23r* induction in response to db-cAMP, IL-23 and in combination (Fig. 3, F and G).

These results together suggest that the PGE₂-EP2/EP4-cAMP-PKA signaling works together with IL-23 signaling to activate STAT3, CREB1 and NF- κ B for induction of *Il23r* expression in Th17 cells.

Gene signature induced by PGE₂-EP2/EP4-cAMP signaling in CD4⁺ T cell populations primed with IL-6 and TGF- β 1

Since pathogenic Th17 cells should express various molecules in addition to IL-23R to exert their pathogenicity, we next examined how PGE₂-EP2/EP4-cAMP signaling contributes to expression of such pathogenic genes in Th17 cells. CD4⁺ T cells were cultured under the Th17-skewing conditions with IL-6 and TGF- β 1 for 3 days, then incubated with db-cAMP alone, IL-23 alone or both for 24 h, and subjected to microarray analysis. The numbers of genes up/down-regulated more than 2-folds by each stimulation were examined by Venn-diagrams (Fig. 4, A), and the genes expressed in each cluster (Table E, 1-8) were subjected to heat-map analysis (Fig E3, A) and gene ontology analysis (Fig E3, B; Table E, 9-11). Expression of representative genes in each

cluster is shown in the heat-map (Fig. 4, B). Cluster 1U included genes (e.g. *Il17a*,
Il17f, *Il1r1* and *Il23r*) that were up-regulated by db-cAMP, IL-23, or both in
combination (Fig. 4 B, left). Cluster 2U included genes (e.g. *Il22*) that were increased
by IL-23 alone or its combination with db-cAMP (Fig. 4 B, left). Cluster 3U
encompasses various genes, which up-regulated by db-cAMP alone or its combination
with IL-23 but not IL-23 alone. They include genes involved in cell migration and
adhesion such as *Ccr2*, *Cxcr4*, *Cx3cr1*, *Ccr6*, *Slpr1*, *Sema4f*, *Sema6c*, *Efna2*, *Sell*, *Selp*
and *Itgb3*, those involved in induction of IFN- γ , such as *Il12rb2*, *Il18r1* and *Il18rap*,
and those involved in cell activation such as *Tlr4*, *Tgfb3*, *Rasa*, *Rasgrp2*, *Lat2*, *Txk* and
Rora (Fig. 4 B, left). Cluster 4U include genes such as *Il1b*, *Il17rc*, *Il17re*, *Prkcq*,
Sema3c, *Sema6a* and *Tlr12* that are up-regulated by the combination of IL-23 and db-
cAMP only (Fig. 4 B, left). On the other hand, the genes in Cluster 3D and Cluster 4D
were down-regulated by db-cAMP, and contained *Il10*, *Il2*, *Il4*, *Il5*, *Il13* and *Il9*, which
are known as suppressive factors of inflammation, (Fig. 4 B, right). Expression of the
representative genes was then confirmed by qRT-PCR analysis. Expression of *Il17a*,
Il17f and *Il23r* in Cluster 1U, *Il18r1*, *Il18rap*, *Slpr1*, *Ccr2*, *Cxcr4*, *Tlr4*, *Cxcl3*, *Cx3cr1*,
Sema4f, *Sell* and *Txk* in Cluster 3U and *Il17re*, *Sema3c* and *Sema6a* in Cluster 4U was
all up-regulated (Fig. 4, C), and expression of *Il10* in Cluster 3D was down-regulated by
the addition of db-cAMP compared to incubation with IL-23 alone (Fig. 4, D). Thus,
signaling through cAMP regulates expression of various genes that are not regulated by
IL-23 alone, and may confer pathogenic property to Th17 cells.

**T cell-intrinsic PGE₂-EP2/EP4 signaling is critical in IL-23-mediated psoriatic skin
inflammation *in vivo***

Accumulating evidences suggest that Th17 cells become pathogenic via the IL-23-IL23 receptor axis and play crucial roles in development of various autoimmune diseases including psoriasis.^{8,56,57} However, how these Th17 cells acquired the pathogenicity *in vivo* and to what extent the microenvironment of diseases contributes to this process remain to be defined. In the IL-23-induced psoriasis mouse model, gene expression of enzymes involved in PGE₂ biosynthesis including *Ptgs2* encoding COX-2, *Ptges* encoding membrane-associated PGE synthase, mPGES1, and *Ptges2* encoding membrane-associated PGE synthase-2, mPGES2, were all up-regulated by IL-23 administration into the skin (Fig E4, A), which is consistent with clinical observation that local PGE₂ levels are elevated in blister fluids from human psoriatic skin.⁵⁸ We therefore hypothesized that IL-23 possibly activates PGE₂–EP2/EP4 signaling, which may contribute to psoriasis pathogenesis.

To test this hypothesis, we injected IL-23 into the skin of WT C57BL/6N mice as well as EP2 knockout (KO) mice⁴⁶ with or without administration with a EP4 antagonist, AS1954813,⁴⁹ and assessed skin inflammation by ear thickness and histology. The EP2 deficiency or the EP4 antagonism alone reduced IL-23-induced ear swelling by half and attenuated edema and cell infiltration, and, when combined, led to nearly complete suppression of IL-23-dependent skin inflammation (Fig. 5, A and B). Blockade of EP2 and/or EP4 caused no alteration in PBS-injected control ear (Fig E3, B). To examine at which step of inflammation EP2 deficiency and EP4 antagonism exert their effects and whether it is related to generation of pathogenic Th17 cells, we digested ear tissues and analyzed CD4⁺ T cell populations in the skin by flow cytometry. While there were little numbers of cells producing IL-17A or IFN- γ in PBS-injected control ear, significant accumulation of the IL-17A⁺ and IL-17A⁺IFN- γ ⁺ CD4⁺ T cell

populations were observed in the IL-23-injected ear as observed in psoriasis dermis in psoriasis patients.⁵⁹ The IL-17A⁺IFN- γ ⁺ CD4⁺ T cell population is suggested one population of pathogenic Th17 cells.⁶⁰ This CD4⁺ T cell population was shown to arise in an IL-23-dependent manner from adoptively transferred T cells in transfer colitis,²⁶ and may reflect the Th17 to Th1 reprogramming at inflammatory sites as shown for antigen-specific Th17 cells transferred to NOD mice.²² This accumulation was significantly reduced by blockade of either EP2 or EP4 alone and nearly completely suppressed by blockade of both EP2 and EP4 (Fig. 5, C and Fig E4, C-E). Consistently, expression of *Il17a* and *Ifng* that was up-regulated in the IL-23-injected ear was also reduced to the negligible levels by combined EP2 and EP4 blockade (Fig. 5 D, left and middle). Notably, EP2 and EP4 blockade also markedly inhibited enhanced expression of *Il23r* by IL-23 injection (Fig. 5 D, right). These findings together indicate that the EP2/EP4 signaling is indeed involved in the generation of pathogenic Th17 cells and elicitation of inflammation in this model. We then asked whether T-cell intrinsic EP2/EP4 signaling is responsible for these IL-23-induced phenotypes. To this end, we used EP2^{flox/flox} mice⁴⁷ and EP4^{flox/flox} mice⁴⁸ and generated EP2^{flox/flox} EP4^{flox/flox} Lck-Cre⁺ mice. EP2^{flox/flox} EP4^{flox/flox} Lck-Cre⁺ mice showed no significant differences in the numbers of total cells, B cells, T cells, CD4 T cells, CD8 T cells, Th1 cells, Th17 cells and Treg cells in the thymus, spleen, lymph node and peripheral blood compared to control WT Lck-Cre⁺ mice (Fig E5, A). However, deficiency of both EP2 and EP4 selectively in T cells prevented accumulation of Th17 cells in the ear and almost completely attenuated IL-23-induced skin inflammation (Fig. 5, E and F). These results together therefore suggest that the T cell-intrinsic PGE₂-EP2/EP4 signaling is critical for the generation of pathogenic Th17 cells in psoriasis model. We also performed

imiquimod (IMQ)-induced psoriasis model⁸, in which we applied IMQ to the ear of WT or EP2 KO mice with or without EP4 antagonist for 6 days (Fig E6, A). We found that ear swelling was also significantly reduced by EP2 deficiency and EP4 antagonism and additively in combination similar to the results in IL-23-induced psoriasis model.

Given the above findings, we next examined the effects of COX inhibitors on skin inflammation in IL-23-induced psoriasis model (Fig E6, B and C). Treatment with indomethacin and SC-236 significantly suppressed the IL-23-induced ear swelling with concomitant suppression of IL-17A⁺ and IL-17A⁺ IFN- γ ⁺ cells in the skin (Fig E6, B and C). These findings together suggest that COX inhibitors are as potent as EP2 and EP4 antagonists in suppressing skin inflammation at least in this model.

PGE₂ signaling positively correlates with the IL-23/Th17 pathway in human psoriatic skin biopsies

Finally, to extrapolate our findings in mice to humans, we analyzed a public microarray dataset on gene expression profiles in skin biopsies from psoriasis patients and healthy control individuals,⁶¹ with a particular interest in correlation of PGE₂ signaling and the IL-23/Th17 pathway. As expected, psoriatic lesional skin overexpressed Th17 signature genes (including *IL23A*, *IL12B*, *IL23R*, *IL17A*, *IL17F*, and *IL22*), *STAT3* and *NFKB1* (encoding NF- κ B p105) (Fig. 6, A). Moreover, psoriatic lesional skin overexpressed enzymes in PGE₂ biosynthesis, e.g., *PTGS2*, *PTGES* and *PTGES2*, and the EP4 receptor (*PTGER4*) but under-expressed the PGE₂ degrading enzyme, 15-PGDH (encoded by *HPGD*) (Fig. 6, A). Interestingly, expression of Th17 signature genes positively correlated with those involved in PGE₂ biosynthesis (e.g. *PTGES* and *PTGES2*) and receptor (e.g. *PTGER4*) but negatively correlated with *HPGD* (Fig. 6, B). In addition,

400 the clinically effective anti-IL-23 therapy⁶² down-regulated gene expression of not only
401 the IL-23/IL-17 pathway (e.g., *IL23A*, *IL23R*, *IL17A*) but also those in PGE synthesis
402 like *PTGES* (Fig. 6, C and D). These findings support a potential crosstalk between the
403 PGE₂ and IL-23/IL-17 pathways also in human psoriatic skin inflammation.

DISCUSSION

The IL-23-IL-23 receptor signaling plays a critical role in generation of pathogenic Th17 cells in autoimmunity.⁵⁻⁹ However, there remain several issues to be solved on this action, namely how this signaling gets promoted, what transcriptional mechanisms other than STAT3 are involved, what, along with the IL-23 signaling, makes Th17 cells pathogenic, whether and how much such mechanism operates *in vivo* and how relevant are the findings obtained in mouse to humans. Given the previously reported action of PGE₂ on Th17 expansion,⁴³⁻⁴⁵ we focused here on PGE₂ action in Th17 pathogenicity to address these issues.

We have first found that PGE₂ synergizes with IL-23 and enhances *Il23r* expression through EP2 and EP4, a finding consistent with the findings in human Th17 cells.⁴⁴ We have then found that IL-23 stimulation induces PGE₂ production in Th17 cells and the IL-23-induced *Il23r* expression was attenuated by the treatment of cells with indomethacin or EP2/EP4 antagonists. These results thus suggest a previously unsuspected intrinsic amplification mechanism mediated by the PGE₂-EP2/EP4 signaling in Th17 cells that helps trigger the initial IL-23 responses in premature Th17 cells.

We have further analyzed the transcriptional mechanisms underlying the synergistic action of IL-23 and PGE₂, and found that this action is mediated by not only STAT3 but also CREB1 and NF- κ B. The involvement of CREB1 is analogous to that in the PGE₂-EP2/EP4-mediated *Il12rb2* induction during Th1 cell differentiation,⁴² and may be consistent with the findings by Hernandez *et al.*⁶³ showing that the CREB1/CRTC2 pathway regulates expression of IL-17A and IL-17F and that Th17 differentiation is defective in CRTC2 mutant mice. IL-23R and IL-12R β 2 make a pair

with the same molecule, IL-12R β 1, to form IL-23 receptor and IL-12 receptor, respectively. It is interesting that the same pathway regulates expression of these two genes. We have also used T cells from p105 NF- κ B1-deficient mice and CTP-NBD and unraveled the involvement of NF- κ B in the IL-23/cAMP-induced *Il23r* expression in Th17 cells. Consistent with these findings, we previously found that PGE₂ through EP2 or EP4 activates NF- κ B1-containing NF- κ B in various types of cells including macrophages and dendritic cells, and induces expression of inflammation-related genes including COX-2, which then produces PGE₂ and amplifies this process.^{47,54,64} Our present findings thus further extend the importance of this COX-2-PGE₂-EP2/EP4-NF- κ B loop to generation of Th17 cell pathogenicity. On the other hand, Boniface *et al.* suggested that PGE₂-induced enhancement of *Il23R* expression in human Th17 cells was mediated by IL-1 β -IL-1 receptor pathway.⁴⁴ This is also a possibility in mice, because up-regulated expression of *Il1r1* and *Il1b* was detected in Cluster 1U and Cluster 4U by our microarray analysis (Fig. 4 B, left). However, we assume that this mechanism does not critically operate in our experiment, because the addition of anti-IL-1 β antibody to the medium did not reduce the *Il23r* induction (Fig E7).

In addition to *Il23r*, our microarray analysis has revealed that stimulation of the EP2/EP4 signaling together with IL-23 facilitates expression of a variety of pathogenic Th17 signature genes (i.e. *Il17a*, *Il17f*, *Il18r1* and *Tgfb3*). Interestingly, PGE₂-EP2/EP4 signaling also up-regulated the expression of various genes related to chemotaxis and migration such as *Slpr1*, *Ccr2*, *Cxcl3*, *Cx3cr1*, *Cxcr4*, *Sema4f*, *Sell*, *Sema3c* and *Sema6a* (Fig. 4 B, left). These results suggest that PGE₂-EP2/EP4 signaling may contribute to migration, infiltration and accumulation of Th17 cells into inflammation

lesion. On the other hand, the addition of db-cAMP down-regulated expression of *Il10*, *Il2*, *Il4* and *Il9*, which are known as suppressive factors for Th17 cells. Although some of these results such as IL-17A are consistent with the previous findings in human Th17 cells,⁴⁴ our study did not detect induction of IFN- γ and T-bet in cultured Th17 cells, which may reflect the stages of Th17 cells examined in each study.^{20,24,65} It should also be noted that our analysis was carried out in the whole CD4⁺ T cell population pretreated with IL-6 and TGF- β 1 and stimulated with each stimulus, in which IL-17A⁺ cells comprise about 10%. Single cell RNA sequencing analysis is therefore desired to establish the gene expression signatures specific to Th17 cells matured with each stimulus.

Nonetheless, the most important point in our study was that the EP2/EP4 signaling in Th17 cells identified here is critical in eliciting their pathogenicity in vivo in immune inflammation. We tested this issue in IL-23-induced mouse psoriasis model. Intriguingly, not only the systemic inhibition of EP2/EP4 signaling using the EP4 antagonist in EP2 KO mice but also selective loss of EP2 and EP4 in T cells almost completely suppressed inflammation induced by IL-23. This was accompanied by suppression of accumulation of IL-17A⁺ and IL-17A⁺IFN- γ ⁺ T cells and suppression of expression of *Il17a*, *Ifng*, *Il23r* genes in the lesion. These results suggest that the PGE₂-EP2/EP4 signaling functions critically in generation of pathogenic Th17 cells induced by IL-23 *in situ*. Of those Th17 cells, antigen-specific Th17 cells were shown to be specifically involved in the pathogenesis of mouse models of autoimmune inflammation including EAE,⁶⁶ type 1 diabetes,²² and psoriasis.⁶⁷ Quite recently it was also reported that mPGES1 is involved in generation of antigen-specific Th17 cells by regulating PGE₂ production in a T cell-autocrine and paracrine manner.⁶⁸ Our present findings

combined together with these findings suggest that PGE₂ plays an important role in psoriasis through the regulation of antigen-specific pathogenic Th17 cells.

The present study also showed that EP2 deficiency and EP4 antagonism also significantly suppressed the psoriatic inflammation in IMQ model. Notably, however, the combined EP2 deficiency and EP4 antagonism did not completely suppress ear swelling in this model, possibly because there is the IL-17-independent component in skin inflammation in this model.⁸

In this study, we also tested the effect of COX inhibitors in IL-23-induced psoriasis model, and found that COX inhibitors are as potent as EP2 and EP4 antagonist in suppressing psoriasis-like skin inflammation in this model. The question is whether COX inhibitors are beneficial in Th17-driven human autoimmune diseases. COX inhibitors, particularly Celecoxib, are used for treatment of the early stage of rheumatoid arthritis patients, and in patients with mild psoriatic arthritis.⁶⁹ In these cases, COX inhibitors produce good symptomatic relief. While this effect is ascribed to their analgesic and general anti-inflammatory actions, our study suggests that it may be derived at least in part from their suppressive action on Th17-mediated pathology, a possibility that should be tested in future. On the other hand, COX inhibitors have less appreciable therapeutic benefits in established psoriasis and advanced RA in human patients. There are several plausible reasons. PG-mediated process may be critical in triggering pathogenic Th17 cell generation, but not so in advanced stage of diseases that might be dominantly regulated by established Th17 cells. Another may be due to the fact that PGs cause immune inflammation not by acting alone but by working with cytokines and boosting and modifying their actions. COX inhibitors may therefore exert therapeutic benefits more effectively when combined with anti-cytokine drugs, and

lessen the dose of the latter. Finally, COX inhibitors may divert arachidonate metabolism to leukotriene. Recent studies suggest that leukotrienes facilitate maturation and migration of Th17 cells.^{70,71} Further studies need to be conducted to unravel these issues.

Another topic to be discussed on PGE₂ in psoriasis is its facilitative action in ultraviolet (UV) irradiation therapy, which at a glance contradicts our present findings on the facilitative action of PGE₂ on Th17 pathogenicity. UVB-irradiation is an effective therapeutic treatment of psoriasis by inducing immunosuppression.⁷² We previously showed that UVB induces PGE₂ in the epidermis and PGE₂-EP4 signaling mediates systemic immunosuppression via the up-regulation of RANKL in keratinocytes and inducing regulatory T cells.⁷³ Thus, the PGE₂-EP4 signaling in this case facilitates immunosuppression and not immune activation. One point is that UVB does not penetrate to the dermis and the events it causes are within the epidermis, while IL-23-induced inflammatory events occur in the dermis. Another point is difference in the context, UVB irradiation in the UV therapy and IL-23 in psoriatic inflammation. PGE₂ alone does not induce either effects, but functions directionally dependent on the context.

Finally, we have examined the relevance of our findings to human diseases by analyzing biopsy samples from psoriasis patients. Psoriatic lesional skin over-expressed not only Th17 signature genes including *IL23A*, *IL12B*, *IL23R*, *IL17A*, *IL17F*, *IL22*, *STAT3* and *NFKB1*, but also those involved in PGE₂ biosynthesis and function such as *PTGS2*, *PTGES*, *PTGES2* and *PTGER4*. Expression of Th17 signature genes shows positive correlation with *PTGES*, *PTGES2* and *PTGER4*, and negative correlation with *HPGD*, and the anti-IL-23 therapy down-regulated expression of not only genes in the

IL-23/IL-17 pathway (e.g., *IL23A*, *IL23R*, *IL17A*) but also those in PGE₂ synthesis, suggesting that these two are functionally linked. These findings together with the finding by Kofler *et al.* that EP2 is expressed in Th17 cells from multiple sclerosis patients and that forced expression of EP2 in healthy Th17 cells triggers expression of pathogenic genes⁷⁴ indicate that T cell-intrinsic EP2/EP4 signaling is critical in IL-23-driven Th17 cell pathogenesis also in humans, and support a view that the combined inhibition of EP2 and EP4 is of value in therapeutic intervention of various Th17-mediated diseases.

ACKNOWLEDGMENTS

We thank A. Kakizuka for suggestions and encouragements, T. Arai for secretarial assistance and K. Naruo for assistance with animal experiments.

REFERENCES

1. Dong C. TH17 cells in development: an updated view of their molecular identity and genetic programming. *Nat Rev Immunol* 2008;8:337-348. <https://doi.org/10.1038/nri2295>
2. Kemper C, Atkinson JP. T-cell regulation: with complements from innate immunity. *Nat Rev Immunol* 2007;7:9-18. <https://doi.org/10.1038/nri1994>
3. Korn T, Bettelli E, Oukka M, Kuchroo VK. IL-17 and Th17 Cells. *Annu Rev Immunol* 2009;27:485-517. <https://doi.org/10.1146/annurev.immunol.021908.132710>

- 547
- 548 4. Zhu J, Paul WE. Peripheral CD4⁺ T-cell differentiation regulated by networks of
- 549 cytokines and transcription factors. *Immunol Rev* 2010;238:247-262.
- 550 <https://doi.org/10.1111/j.1600-065X.2010.00951.x>
- 551
- 552 5. Cua DJ, Tato CM. Innate IL-17-producing cells: the sentinels of the immune system.
- 553 *Nat Rev Immunol* 2010;10:479-489. <https://doi.org/10.1038/nri2800>
- 554
- 555 6. Murphy CA, Langrish CL, Chen Y, Blumenschein W, McClanahan T, Kastelein RA,
- 556 et al. Divergent pro- and antiinflammatory roles for IL-23 and IL-12 in joint
- 557 autoimmune inflammation. *J Exp Med* 2003;198:1951-1957.
- 558 <https://doi.org/10.1084/jem.20030896>
- 559
- 560 7. Torchinsky MB, Blander JM. T helper 17 cells: discovery, function, and physiological
- 561 trigger. *Cell Mol Life Sci* 2010;67:1407-1421. [https://doi.org/10.1007/s00018-009-](https://doi.org/10.1007/s00018-009-0248-3)
- 562 [0248-3](https://doi.org/10.1007/s00018-009-0248-3)
- 563
- 564 8. van der Fits L, Mourits S, Voerman JS, Kant M, Boon L, Laman JD, et al.
- 565 Imiquimod-induced psoriasis-like skin inflammation in mice is mediated via the IL-
- 566 23/IL-17 axis. *J Immunol* 2009;182:5836-5845.
- 567 <https://doi.org/10.4049/jimmunol.0802999>
- 568
- 569 9. Yen D, Cheung J, Scheerens H, Poulet F, McClanahan T, McKenzie B, et al. IL-23 is
- 570 essential for T cell-mediated colitis and promotes inflammation via IL-17 and IL-6. *J*

Clin Invest 2006;116:1310-1316. <https://doi.org/10.1172/JCI21404>

10. Kopp T, Riedl E, Bangert C, Bowman EP, Greisenegger E, Horowitz A, et al. Clinical improvement in psoriasis with specific targeting of interleukin-23. *Nature* 2015;521:222-226. <https://doi.org/10.1038/nature14175>

11. Krueger JG, Ferris LK, Menter A, Wagner F, White A, Visvanathan S, et al. Anti-IL-23A mAb BI 655066 for treatment of moderate-to-severe psoriasis: Safety, efficacy, pharmacokinetics, and biomarker results of a single-rising-dose, randomized, double-blind, placebo-controlled trial. *J Allergy Clin Immunol* 2015;136:116-124. <https://doi.org/10.1016/j.jaci.2015.01.018>

12. Leonardi CL, Kimball AB, Papp KA, Yeilding N, Guzzo C, Wang Y, et al. Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 76-week results from a randomised, double-blind, placebo-controlled trial (PHOENIX 1). *Lancet* 2008;371:1665-1674. [https://doi.org/10.1016/S0140-6736\(08\)60725-4](https://doi.org/10.1016/S0140-6736(08)60725-4)

13. Papp KA, Langley RG, Lebwohl M, Krueger GG, Szapary P, Yeilding N, et al. Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 52-week results from a randomised, double-blind, placebo-controlled trial (PHOENIX 2). *Lancet* 2008;371:1675-1684. [https://doi.org/10.1016/S0140-6736\(08\)60726-6](https://doi.org/10.1016/S0140-6736(08)60726-6)

14. Teng MW, Bowman EP, McElwee JJ, Smyth MJ, Casanova JL, Cooper AM, et al. IL-12 and IL-23 cytokines: from discovery to targeted therapies for immune-mediated inflammatory diseases. *Nat Med* 2015;21:719-729. <https://doi.org/10.1038/nm.3895>
15. Ivanov II, McKenzie BS, Zhou L, Tadokoro CE, Lepelley A, Lafaille JJ, et al. The orphan nuclear receptor ROR γ directs the differentiation program of proinflammatory IL-17+ T helper cells. *Cell* 2006;126:1121-1133. <https://doi.org/10.1016/j.cell.2006.07.035>
16. Morishima N, Mizoguchi I, Takeda K, Mizuguchi J, Yoshimoto Y. TGF-beta is necessary for induction of IL-23R and Th17 differentiation by IL-6 and IL-23. *Biochem Biophys Res Commun* 2009;386:105-110. <https://doi.org/10.1016/j.bbrc.2009.05.140>
17. Yang XO, Pappu BP, Nurieva R, Akimzhanov A, Kang HS, Chung Y, et al. T helper 17 lineage differentiation is programmed by orphan nuclear receptors ROR alpha and ROR gamma. *Immunity* 2008;28:29-39. <https://doi.org/10.1016/j.immuni.2007.11.016>
18. Zhou L, Ivanov II, Spolski R, Min R, Shenderov K, Egawa T, et al. IL-6 programs T(H)-17 cell differentiation by promoting sequential engagement of the IL-21 and IL-23 pathways. *Nat Immunol* 2007;8:967-974. <https://doi.org/10.1038/ni1488>
19. Zhou L, Lopes JE, Chong MM, Ivanov II, Min R, Victora GD, et al. TGF-beta-induced Foxp3 inhibits T(H)17 cell differentiation by antagonizing ROR γ

function. *Nature* 2008;453:236-240. <https://doi.org/10.1038/nature06878>

20. Lee Y, Awasthi A, Yosef N, Quintana FJ, Xiao S, Peters A, et al. Induction and molecular signature of pathogenic TH17 cells. *Nat Immunol* 2012;13:991-999. <https://doi.org/10.1038/ni.2416>

21. Gagliani N, Amezcu Vesely MC, Iseppon A, Brockmann L, Xu H, Palm NW, et al. Th17 cells transdifferentiate into regulatory T cells during resolution of inflammation. *Nature* 2015;523:221-225. <https://dx.doi.org/10.1038/nature14452>

22. Li CR, Mueller EE, and Bradley LM. Islet antigen-specific Th17 cells can induce TNF- α dependent autoimmune diabetes. *J Immunol* 2014;192:1425-32. <https://dx.doi.org/10.4049/jimmunol.1301742>

23. Stockinger B and Omenetti S. The dichotomous nature of T helper 17 cells. *Nat Rev Immunol* 2017;17:535-544. <https://dx.doi.org/10.1038/nri.2017.50>

24. Wu C, Yosef N, Thalhamer T, Zhu C, Xiao S, Kishi Y, et al. Induction of pathogenic TH17 cells by inducible salt-sensing kinase SGK1. *Nature* 2013;496:513-517. <https://doi.org/10.1038/nature11984>

25. Horste GM, Wu C, Wang C, Cong L, Pawlak M, Lee Y, et al. RBPJ controls development of pathogenic Th17 cells by regulating IL-23 receptor expression. *Cell Reports* 2016;16:392-404. <https://dx.doi.org/10.1016/j.celrep.2016.05.088>

26. Ahern PP, Schiering C, Buonocore S, McGeachy MJ, Cua DJ, Maloy KJ, et al. Interleukin-23 drives intestinal inflammation through direct activity on T cells. *Immunity* 2010;33:279-288. <https://dx.doi.org/10.1016/j.immuni.2010.08.010>
27. Harbour SN, Maynard CL, Zindl CL, Schoeb TR, and Weaver CT. Th17 cells give rise to Th1 cells that are required for the pathogenesis of colitis. *PNAS* 2015;112:7061-7066. <https://dx.doi.org/10.1073/pnas.1415675112>
28. Anderson CA, Boucher G, Lees CW, Franke A, D'Amato M, Taylor KD, et al. Meta-analysis identifies 29 additional ulcerative colitis risk loci, increasing the number of confirmed associations to 47. *Nat Genet* 2011;43:246-252. <https://dx.doi.org/10.1038/ng.764>
29. Franke A, McGovern DP, Barrett JC, Wang K, Radford-Smith GL, Ahmad T, et al. Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. *Nat Genet* 2010;42:1118-1125. <https://doi.org/10.1038/ng.717>
30. Stuart PE, Nair RP, Tsoi LC, Tejasvi T, Das S, Kang HM, et al. Genome-wide association analysis of psoriatic arthritis and cutaneous psoriasis reveals differences in their genetic architecture. *Am J Hum Genet* 2015;97:816-836. <https://doi.org/10.1016/j.ajhg.2015.10.019>
31. Burkett PR, Meyer zu Horste G, Kuchroo VK. Pouring fuel on the fire: Th17 cells,

the environment, and autoimmunity. *J Clin Invest* 2015;125:2211-2219.

<https://doi.org/10.1172/JCI78085>

32. Gaffen SL, Jain R, Garg AV, Cua DJ. The IL-23-IL-17 immune axis: from mechanisms to therapeutic testing. *Nat Rev Immunol* 2014;14:585-600.

<https://doi.org/10.1038/nri3707>

33. Boyman O, Comte D, Spertini F. Adverse reactions to biologic agents and their medical management. *Nat Rev Rheumatol* 2014;10:612-27.

<https://doi.org/10.1038/nrrheum.2014.123>

34. Hawkes JE, Chan TC, Krueger JG. Psoriasis pathogenesis and the development of novel targeted immune therapies. *J Allergy Clin Immunol* 2017;140:645-653.

<https://doi.org/10.1016/j.jaci.2017.07.004>

35. Sandborn WJ, Ghosh S, Panes J, Vranic I, Su C, Rousell S, et al. tofacitinib, an oral janus kinase inhibitor, in active ulcerative colitis. *N Engl J Med* 2012;367:616-624.

<http://doi.org/10.1056/NEJMoa1112168>

36. Hirata T, Narumiya S. Prostanoid receptors. *Chem Rev* 2011;111:6209-6230.

<https://doi.org/10.1021/cr200010h>

37. Harris SG, Padilla J, Koumas L, Ray D, Phipps RP. Prostaglandins as modulators of immunity. *Trends Immunol* 2002;23:144-150.

38. Sreeramkumar V, Fresno M, Cuesta N. Prostaglandin E2 and T cells: friends or foes? *Immunol Cell Biol* 2012;90:579-586. <https://doi.org/10.1038/icb.2011.75>
39. Aoki T, Narumiya S. Prostaglandins and chronic inflammation. *Trends Pharmacol Sci* 2012;33:304-311. <https://doi.org/10.1016/j.tips.2012.02.004>
40. Hirata T, Narumiya S. Prostanoids as regulators of innate and adaptive immunity. *Adv Immunol* 2012;116:143-174. <https://doi.org/10.1016/B978-0-12-394300-2.00005-3>
41. Sakata D, Yao C, Narumiya S. Prostaglandin E2, an immunoactivator. *J Pharmacol Sci* 2010;112:1-5.
42. Yao C, Hirata T, Soontrapa K, Ma X, Takemori H, Narumiya S. Prostaglandin E(2) promotes Th1 differentiation via synergistic amplification of IL-12 signalling by cAMP and PI3-kinase. *Nat Commun* 2013;4:1685. <https://doi.org/10.1038/ncomms2684>
43. Yao C, Sakata D, Esaki Y, Li Y, Matsuoka T, Kuroiwa K, et al. Prostaglandin E2-EP4 signaling promotes immune inflammation through Th1 cell differentiation and Th17 cell expansion. *Nat Med* 2009;15:633-640. <https://doi.org/10.1038/nm.1968>
44. Boniface K, Bak-Jensen KS, Li Y, Blumenschein WM, McGeachy MJ, McClanahan

- TK, et al. Prostaglandin E2 regulates Th17 cell differentiation and function through cyclic AMP and EP2/EP4 receptor signaling. *J Exp Med* 2009;206:535-548. <https://doi.org/10.1084/jem.20082293>
45. Napolitani G, Acosta-Rodriguez EV, Lanzavecchia A, Sallusto F. Prostaglandin E2 enhances Th17 responses via modulation of IL-17 and IFN-gamma production by memory CD4+ T cells. *Eur J Immunol* 2009;39:1301-1312. <https://doi.org/10.1002/eji.200838969>
46. Hizaki H, Segi E, Sugimoto Y, Hirose M, Saji T, Ushikubi F, et al. Abortive expansion of the cumulus and impaired fertility in mice lacking the prostaglandin E receptor subtype EP(2). *Proc Natl Acad Sci U S A* 1999;96:10501-10506.
47. Aoki T, Frosen J, Fukuda M, Bando K, Shioi G, Tsuji K, et al. Prostaglandin E2-EP2-NF-kappaB signaling in macrophages as a potential therapeutic target for intracranial aneurysms. *Sci Signal* 2017;10: eaah6037. <https://doi.org/10.1126/scisignal.aah6037>
48. Schneider A, Guan Y, Zhang Y, Magnuson MA, Pettepher C, Loftin CD, et al. Generation of a conditional allele of the mouse prostaglandin EP4 receptor. *Genesis* 2004;40:7-14. <https://doi.org/10.1002/gene.20048>
49. Zenkoh T, Nozawa E, Matsuura M, Seo R. Ornithine derivative. In Google Patents. 2008.

50. Ohno K, Fujiwara M, Fukushima M, Narumiya S. Metabolic dehydration of prostaglandin E2 and cellular uptake of the dehydration product: correlation with prostaglandin E2-induced growth inhibition. *Biochem Biophys Res Commun* 1986;139:808-15.
51. Aoki T, Nishimura M, Matsuoka T, Yamamoto K, Furuyashiki T, Kataoka H, et al. PGE2-EP2 signalling in endothelium is activated by haemodynamic stress and induces cerebral aneurysm through an amplifying loop via NF- κ B. *Br J Pharmacol* 2011; 163: 1237–1249. <http://doi.org/10.1111/j.1476-5381.2011.01358.x>
52. Che Mat NF, Zhang X, Guzzo C, Gee K. Interleukin-23-induced interleukin-23 receptor subunit expression is mediated by the Janus kinase/signal transducer and activation of transcription pathway in human CD4 T cells. *J Interferon Cytokine Res* 2011;31:363-371. <https://doi.org/10.1089/jir.2010.0083>
53. Best JL, Amezcua CA, Mayr B, Flechner L, Murawsky CM, Emerson B, et al. Identification of small-molecule antagonists that inhibit an activator: coactivator interaction. *Proc Natl Acad Sci U S A* 2004;101:17622-17627. <https://doi.org/10.1073/pnas.0406374101>
54. Ma X, Aoki T, Narumiya S. Prostaglandin E2-EP4 signaling persistently amplifies CD40-mediated induction of IL-23 p19 expression through canonical and non-canonical NF- κ B pathways. *Cell Mol Immunol* 2016;13:240-250.

<https://doi.org/10.1038/cmi.2015.70>

55. Christian F, Smith EL, Carmody RJ. The regulation of NF-kappaB subunits by phosphorylation. *Cells* 2016;5:12. <https://doi.org/10.3390/cells5010012>

56. Lowes MA, Russell CB, Martin DA, Towne JE, Krueger JG. The IL-23/T17 pathogenic axis in psoriasis is amplified by keratinocyte responses. *Trends Immunol* 2013;34:174-181. <https://doi.org/10.1016/j.it.2012.11.005>

57. Rizzo HL, Kagami S, Phillips KG, Kurtz SE, Jacques SL, Blauvelt A. IL-23-mediated psoriasis-like epidermal hyperplasia is dependent on IL-17A. *J Immunol* 2011;186:1495-1502. <https://doi.org/10.4049/jimmunol.1001001>

58. Reilly DM, Parslew R, Sharpe GR, Powell S, Green MR. Inflammatory mediators in normal, sensitive and diseased skin types. *Acta Derm Venereol* 2000;80:171-174.

59. Lowes MA, Kikuchi T, Fuentes-Duculan J, Cardinale I, Zaba LC, Haider AS, et al. Psoriasis vulgaris lesions contain discrete populations of Th1 and Th17 T cells. *J Invest Dermatol* 2008;128:1207-11. <https://doi.org/10.1038/sj.jid.5701213>

60. Duhon R1, Glatigny S, Arbelaez CA, Blair TC, Oukka M, Bettelli E. Pathogenicity of IFN- γ -producing Th17 cells is independent of T-bet. *J Immunol* 2013;190:4478-82. <http://doi.org/10.4049/jimmunol.1203172>

61. Nair RP, Duffin KC, Helms C, Ding J, Stuart PE, Goldgar D, et al. Genome-wide scan reveals association of psoriasis with IL-23 and NF-kappaB pathways. *Nat Genet* 2009;41:199-204. <https://doi.org/10.1038/ng.311>
62. Sofen H, Smith S, Matheson RT, Leonardi CL, Calderon C, Brodmerkel C, et al. Guselkumab (an IL-23-specific mAb) demonstrates clinical and molecular response in patients with moderate-to-severe psoriasis. *J Allergy Clin Immunol* 2014;133:1032-1040. <https://doi.org/10.1016/j.jaci.2014.01.025>
63. Hernandez JB, Chang C, LeBlanc M, Grimm D, Le Lay J, Kaestner KH, et al. The CREB/CRTC2 pathway modulates autoimmune disease by promoting Th17 differentiation. *Nat Commun* 2015;6:7216. <https://doi.org/10.1038/ncomms8216>
64. Ma X, Aoki T, Tsuruyama T, Narumiya S. Definition of Prostaglandin E2-EP2 Signals in the Colon Tumor Microenvironment That Amplify Inflammation and Tumor Growth. *Cancer Res* 2015;75:2822-2832. <https://doi.org/10.1158/0008-5472.CAN-15-0125>
65. McGeachy MJ, Chen Y, Tato CM, Laurence A, Joyce-Shaikh B, Blumenschein WM, et al. The interleukin 23 receptor is essential for the terminal differentiation of interleukin 17-producing effector T helper cells in vivo. *Nat Immunol* 2009;10:314-324. <https://doi.org/10.1038/ni.1698>
66. Korn T, Mitsdoerffer M, Croxford AL, Awasthi A, Dardalhon VA, Galileos G, et al.

- IL-6 controls Th17 immunity in vivo by inhibiting the conversion of conventional T cells into Foxp3⁺ regulatory T cells. *Proc Natl Acad Sci U S A* 2008;105:18460-5. <http://doi.org/10.1073/pnas.0809850105>
67. Nishimoto S, Kotani H, Tsuruta S, Shimizu N, Ito M, Shichita T, et al. Th17 cells carrying TCR recognizing epidermal autoantigen induce psoriasis-like skin inflammation. *J Immunol* 2013;191:3065-72. <http://doi.org/10.4049/jimmunol.1300348>
68. Maseda D, Johnson EM, Nyhoff LE, Baron B, Kojima F, Wilhelm AJ, et al. mPGES1-dependent prostaglandin E₂ (PGE₂) controls antigen-specific Th17 and Th1 responses by regulating T autocrine and paracrine PGE₂ production. *J Immunol* 2018; 200:725-736. <http://doi.org/10.4049/jimmunol.1601808>
69. Menter A, Gottlieb A, Feldman SR, Van Voorhees AS, Leonardi CL, Gordon KB, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 1. Overview of psoriasis and guidelines of care for the treatment of psoriasis with biologics. *J Am Acad Dermatol* 2008;58:826-50. <http://doi.org/10.1016/j.jaad.2008.02.039>
70. Lee W, Kim HS and Lee GR. Leukotrienes induce the migration of Th17 cells. *Immunology and Cell Biology* 2015;93:472–479. <http://doi.org/10.1038/icb.2014.104>
71. Chen H, Qin J, Wei P, Zhang J, Li Q, Fu L, et al. Effects of leukotriene B₄ and

prostaglandin E2 on the differentiation of murine Foxp3+ T regulatory cells and Th17 cells. Prostaglandins, Leukotrienes and Essential Fatty Acids 2009;80:195-200. <http://doi.org/10.1016/j.plefa.2009.01.006>

72. Nakamura M, Farahnik B, and Bhutani T. Recent advances in phototherapy for psoriasis. F1000 Res 2016;5: 1684. <http://doi.org/10.12688/f1000research.8846.1>

73. Soontrapa K, Honda T, Sakata D, Yao C, Hirata T, Hori S, et al. Prostaglandin E2–prostaglandin E receptor subtype 4 (EP4) signaling mediates UV irradiation-induced systemic immunosuppression. Proc Natl Acad Sci U S A 2011;108:6668-6673. <http://doi.org/10.1073/pnas.1018625108>

74. Kofler DM, Marson A, Dominguez-Villar M, Xiao S, Kuchroo VK, Hafler DA. Decreased RORC-dependent silencing of prostaglandin receptor EP2 induces autoimmune Th17 cells. J Clin Invest 2014;124:2513-2522. <https://doi.org/10.1172/JCI72973>

FIGURE LEGENDS

Figure 1. IL-23 mobilizes the endogenous PGE₂-EP2/EP4-cAMP-PKA pathway to facilitate Th17 expansion through synergistic *Il23r* induction. (A and B) Expansion of Th17 population by PGE₂ and IL-23. CD4⁺ T cells were differentiated with TGF-β1 and IL-6 to Th17 cells for 4 days, then stimulated with 100 nM PGE₂ in the absence or presence of IL-23 (10 ng/ml) for additional 3 days. The cells were examined by FACS for IL-17A and IFN-γ (A) and by qRT-PCR for *Il17a* expression (B). (C-E) Effects of PGE₂, agonists selective to each EP subtype and related compounds on *Il23r* expression. Th17 cells were incubated with 100 nM PGE₂, an agonist selective to each EP subtype, ONO-DI-004 (EP1), ONO-AE1-259 (EP2), ONO-AE-248 (EP3), or ONO-AE1-329 (EP4), 100 μM dibutyl cAMP (db-cAMP), 10 μM forskolin (FSK), 100 μM 3-isobutyl-1-methylxanthine (IBMX) with or without IL-23. *Il23r* expression (C and D) or IL-17A concentrations in culture supernatant (E) was examined. Expression of *Il23r* in Th17 cells stimulated with 100 μM db-cAMP, 300 μM N6-Bnz-cAMP (a PKA agonist), 300 μM 8-pCTP-2'-O-Me-cAMP (an Epac activator) (F), or H-89 (a PKA inhibitor) (G) with or without IL-23. All bars indicate mean ± SEM (n=3, each in A-G). p<0.05, **, p<0.01, ***, p<0.001.

Figure 2. IL-23 self-amplifies its own signaling through a T cell intrinsic positive feedback COX-2-PGE₂-cAMP-IL-23R loop. (A) Expression of COX-2 (*Ptgs2*) mRNA (A) in Th17 cells, or Th17 cells further cultured in the presence or absence of IL-23 for 3 days determined by qRT-PCR. (B) Concentrations of PGE₂ in the culture supernatants of Th17 cells in the presence or absence of IL-23 and indomethacin determined by ELISA. n.d., not detected. (C) *Il23r* expression in Th17 cells stimulated

with PGE₂ and IL-23 in the presence or absence of indomethacin for 3 days. (D) *Il23r* expression in Th17 cells stimulated with PGE₂ and IL-23 in the presence or absence of EP2 (PF-04418948) and/or EP4 (ONO-AE3-208) antagonists for 3 days. All bars indicate mean \pm SEM (n=3, each in A-D). *, p<0.05, **, p<0.01, ***, p<0.001.

Figure 3. STAT3, CREB1 and NF- κ B mediate cAMP and IL-23 induced *Il23r* expression in Th17 cells. (A) Effect of STAT3 inhibitor VII on *Il23r* expression in Th17 cells stimulated with db-cAMP and/or IL-23 for 3 days. (B) Western blot analysis of phospho-Y705 STAT3 and α -Tubulin as a loading control in Th17 cells cultured as described in Supplementary Methods. Representative images from 2 independent experiments are shown. (C) Effect of KG-501 on *Il23r* expression in Th17 stimulated by db-cAMP and/or IL-23. (D) Effects of RNA interference for CREB1 on *Il23r* expression (left) and *Creb1* expression to confirm CREB knockdown efficiency (right). RNA interference, subsequent culture and stimulation of Th17 cells were performed as described in Supplementary Methods. (E) Western blot analysis of phospho-S536 NF- κ B p65 (pp65), phospho-S933 NF- κ B p105 (pp105), p65, p105/p50 and α -Tubulin in Th17 cells stimulated as described in Supplementary Methods. Representative images from 2 independent experiments are shown. (F and G) Effects of p105 KO (F) or CTP-NBD (G) on *Il23r* expression in Th17 cells stimulated with db-cAMP and/or IL-23 for 3 days. All bars indicate mean \pm SEM. (n=3, each in A, C, F, and G, n=18 in B) *, p<0.05, **, p<0.01, ***, p<0.001.

Figure 4. Activation of the COX-2-PGE₂-EP2/EP4-cAMP pathway confers pathogenic Th17 phenotype. (A) Microarray analysis of gene expression profiles in

Th17 cells stimulated with db-cAMP and/or IL-23. Venn-diagram analysis of two folds up- or down-regulated genes compared to the vehicle control upon each stimulus (One-way ANOVA $p < 0.05$, $n = 3$) (left and right, respectively). (B) Heat-map analysis of expression of selected genes from each cluster. (C) qRT-PCR analysis of expression of representative genes of Th17 signature and immune activation in response to db-cAMP, IL-23 or db-cAMP and IL-23 in combination. (D) qRT-PCR analysis of expression of a representative inflammation suppressor gene, *Il10*, in response to db-cAMP, IL-23 or db-cAMP and IL-23 in combination. All bars in C and D indicate mean \pm SEM. ($n = 3$) *, $p < 0.05$, **, $p < 0.01$, ***, $p < 0.001$.

Figure 5. PGE₂-EP2/EP4 signaling in T cells is required for IL-23-driven psoriatic skin inflammation. (A-D) Ear swelling (A) ($n = 16-17$), representative Hematoxylin-Eosin (HE) staining of the histological section of the ear (B) ($n = 3-4$), number of IL-17A⁺ and IL-17A⁺ IFN- γ ⁺ CD4⁺ T cells of the ear (C) and gene expression of *Il17a*, *Ifng* and *Il23r* in the whole ear tissue (D) of WT or EP2 KO mice subcutaneously injected with IL-23 or PBS into the ear daily. An EP4 antagonist (AS1954813, 100 mg/kg) or vehicle was orally administered twice a day to the indicated mice. Bars in (B), 50 μ m. Representative quantification results of the cell number in each population from 4 independent FACS experiments are shown in (C) ($n = 3$). Gene expression was indicated as fold-change compared to PBS-injected ear in (D) ($n = 3$). (E and F) EP2^{flox/flox}EP4^{flox/flox}Lck-Cre⁺ mice and control WT Lck-Cre⁺ mice were subjected to IL-23-induced psoriasis model and the ear swelling (E) ($n = 11$ and 7, respectively) and the number of IL-17A⁺ and IL-17A⁺ IFN- γ ⁺ CD4⁺ T cells in the ear (F) ($n = 7$ and 3, respectively) were analyzed. All bars indicate mean \pm SEM. *, $p < 0.05$, **, $p < 0.01$, ***,

p<0.001.

Figure 6. PGE₂ signaling positively correlates with the IL-23/Th17 pathway in human psoriatic skin biopsies. (A) Expression profiles of genes related to PGE₂ signaling and Th17 signature genes in human non-lesional (NL) or lesional (PL) skin biopsies from patients with psoriasis (n=58) and skin samples from healthy controls (HC, n=64). The z-score transformed values of microarray gene expression dataset GSE13355 were used. Th17 score was generated based on the average expression level of *IL23A*, *IL12B*, *IL23R*, *IL17A*, *IL17F* and *IL22* genes. (B) Correlations of *PTGES*, *PTGES2*, *HPGD* and *PTGER4* gene expression versus that of the Th17 score. Black, green and red dots indicate healthy control, non-lesional and lesional psoriatic biopsies, respectively. (C) Expression profiles of genes related to PGE₂ synthases and Th17 signature genes in human lesional skin biopsies from patients with moderate-to-severe psoriasis before (Baseline, n=22) or 12 weeks after treatment with an IL-23-specific mAb, Guselkumab, (Guselkumab, n=8). The z-score transformed values of microarray gene expression dataset GSE51440 were used. (D) Correlations of gene expression of *PTGS2* and *PTGES* versus that of *IL23R*. P values were calculated by nonparametric Wilcoxon-Mann-Whitney tests (A, C) or nonparametric Spearman correlation test (B, D).




Figure. 1




Figure. 2




Figure. 3




Figure. 4




Figure. 5




Figure. 6

SUPPLEMENTARY METHODS

Purification of CD4⁺ T cells and differentiation into Th17 cells

Spleen was dissected from 6-10 week-old female C57BL/6N mice and cells were dissociated and collected. CD4⁺ T cells were purified from spleen cells by magnetic activated-cell sorting (MACS) using anti-CD4 microbeads (L3T4) (#130-049-201, Miltenyi) on auto-MACS (Miltenyi). The purity of CD4⁺ T cells was ~98 % (n=3) as assessed by FACS (FACS LSR Fortessa, BD Bioscience, San Jose, CA). Purified CD4⁺ T cells were differentiated into Th17 cells by the combination of TGF- β 1 (1 ng/ml, #240-B-002, R&D systems, Minneapolis, MN) and IL-6 (20 ng/ml, R&D systems) in the presence of 5 μ g/ml of anti-CD3 antibody (#14-0031-86, eBioscience, San Diego, CA) and 2.5 μ g/ml of anti-CD28 antibody (#14-0281-86, eBioscience, San Diego, CA) in RPMI-1640 medium containing 10 % fetal bovine serum (FBS) for 4 days. Differentiated cells were then collected, washed and again plated for experiments with TCR stimulation. Experimental condition of each experiment is shown in the **Results** or the **Figure Legends** unless specified otherwise.

17 **Reagents**

18 Agonists selective to each PGE₂ receptor subtype, EP1, EP2, EP3 and EP4
19 (ONO-DI-004, ONO-AE1-259, ONO-AE-248 and ONO-AE1-329 respectively) and an
20 EP4 antagonist, ONO-AE3-208,^{E1} were kindly provided by Ono Pharmaceutical Co.,
21 Osaka, Japan. An EP2 antagonist, PF-04418948, was synthesized according to the
22 previous report.^{E2} An EP4 antagonist, AS1954813, was kindly provided by Astellas
23 Pharmaceutical Co. (Tsukuba, Japan). PGE₂, SC-560 and SC-236 were purchased from
24 Cayman Chemical, Ann Arbor, MI. Dibutyryl cAMP (db-cAMP), forskolin, the
25 N6-Bnz-cAMP, the 8-pCTP-2'-O-Me-cAMP, indomethacin and KG-501 were
26 purchased from Sigma, St. Louis, MO. STAT3 inhibitor VII, Src Kinase Inhibitor-I,
27 H-89, and CPT-NBD peptides were purchased from Calbiochem, San Diego, CA.
28 Mouse IL-1 β /IL-1F2 Antibody (AF-401-SP) was purchased from R&D systems.

29

30 **Quantitative reverse transcription polymerase chain reaction (qRT-PCR)**

31 RNA purification and reverse transcription were performed by the RNeasy Mini Kit
32 (Qiagen GmbH, Hilden, Germany) and the High-capacity cDNA Reverse Transcription
33 Kit (ABI biosystems, Grand Island, NY) according to manufacturers' instructions.
34 cDNA, primers and FastStart DNA MasterPLUS SYBR Green (Takara, Shiga, Japan)
35 were then mixed in 96-well PCR plate, and quantitative PCR was performed using
36 CFX96 Real-Time System (Biorad). The following primers were used in this study;

37 *Gapdh*: forward 5'-TGAACGGGAAGCTCAC-3' and reverse 5'
38 -TCCACCACCCTGTTGC-3'

39 *Il17a*: forward 5'-TGTGAAGGTCAACCTCAAAGTC-3' and reverse 5'
40 -GAGGGATATCTATCAGGGTCTTCA-3'

41 *Il23r*: forward 5'-CCAAGTATATTGTGCATGTGAAGA-3' and reverse 5'
42 -AGCTTGAGGCAAGATATTGTTGT-3'

43 *Ptgs2*: forward 5'- TCGCAGGAAGGGGATGTTGT -3' and reverse 5'-
44 CTGAAGCCCACCCCAAACAC -3'

45 *Creb1*: forward 5'-CCAAACTAGCAGTGGGCAGT-3' and reverse 5'

46 -CCCCATCCGTACCATTGTT-3'

47 *Il17f*: forward 5'-GGAAGACAGCACCATGAAC-3' and reverse

48 5'-TGGACAATGGGCTTGACAG-3'

49 *Il18r1*: forward 5'-GTTGAGATGGAGGATGAGGG-3' and reverse 5'

50 -GACAGAAAACACGCAGGAG-3'

51 *Il18rap*: forward 5'-AGCCTTTAACTCTCCCCTG-3' and reverse 5'

52 -ACACCACCTCTTCCTTCTTC-3'

53 *Slpr1*: forward 5'-CATTCTCATCTGCTGCTTCATC-3' and reverse 5'

54 -CCACAAACATACTCCCTTCCC-3'

55 *Ccr2*: forward 5'-TGAGAAGAAGAGGCACAGG-3' and reverse 5'

56 -CAACAAAGGCATAAATGACAGG-3'

57 *Cxcr4*: forward 5'-ATCTGTGACCGCCTTTACCC-3' and reverse 5'

58 -ATCCTTGCTTGATGACCCCC-3'

- 59 *Tlr4*: forward 5'-CTTTCACCTCTGCCTTCAC-3' and reverse 5'
- 60 -TACAATTCCACCTGCTGCC-3'
- 61 *Cxcl3*: forward 5'-GAACACCCTCAGGCTCAAGG-3' and reverse 5'
- 62 -CCACCAACCAAAGAATACACATGG-3'
- 63 *Cx3cr1*: forward 5'-ACAAAGAGAAAGGACAACGAG-3' and reverse 5'
- 64 -TGATGCGGAAGTAGCAAAAG-3'
- 65 *Sema4f*: forward 5'-AAGAAAGGCAAGAAAGAGGAC-3' and reverse 5'
- 66 -CACATCAATAACCCCGCAC-3'
- 67 *Sell*: forward 5'-TGCCAAGAGACAAACAGAAG-3' and reverse 5'
- 68 -CCAGCCAAATGAGAAATGCC-3'
- 69 *Txk*: forward 5'-CACCGAAAGACATCTCTTCC-3' and reverse 5'
- 70 -ACAACCCCAAACCTGACCAC-3'
- 71 *Il17re*: forward 5'-ACAACCCCAAACCTGACCAC-3' and reverse 5'
- 72 -GGGCAGCAAATCAAAGGAG-3'

73 *Sema3c*: forward 5'-ACAAAGACAGGAGGAAGGAG-3' and reverse 5'-

74 AGTGGCAATGCAGTGGTAG-3'

75 *Sema6a*: forward 5'-GCTCACTCTATGTTGCATTCTC-3' and reverse 5'-

76 ACTTTCCCTTACCCACCCAC-3'

77 *Il10*: forward 5'-TGGGTGAGAAGCTGAAGACC-3' and reverse

78 5'-TTCATGGCCTTGTAGACACC-3'

79 *Ifng*: forward 5'-ATCTGGAGGAACTGGCAAAA-3' and reverse

80 5'-TTCAAGACTTCAAAGAGTCTGAGGTA-3'

81 Expression level of each gene was normalized to that of *Gapdh* and calculated relative

82 to the expression in vehicle-treated group.

83

84 **Measurement of IL-17 and PGE₂ concentration in culture supernatant of Th17**

85 **cells**

86 IL-17 concentration in culture supernatant of differentiated Th17 cells stimulated with

87 100 µM db-cAMP, 10 µM FSK or 100 µM IBMX for 3 days was measured by a Mouse

IL-17 Quantikine ELISA Kit (M1700, R&D systems) according to the manufacturer's instruction.

Th17 cells were stimulated with 10 ng/ml IL-23 for 3 days in the absence or presence of 100 μ M indomethacin and PGE₂ concentration in culture supernatant was determined by a Prostaglandin E₂ ELISA kit - monoclonal (514010, Cayman Chemical, Ann Arbor, MI) according to the manufacturer's instruction.

Gene expression of Th17 cells stimulated IL-23 and/or cAMP from microarray analysis

CD4⁺ T cells were incubated in Th17-skewing condition for 4 days. Differentiated Th17 cells were then stimulated by IL-23, db-cAMP or combination for 24 h. RNA was purified with an RNeasy Mini Kit, amplified and revers transcribed by the High-capacity cDNA Reverse Transcription Kit. cDNA was fragmented and labeled by a Low Input Quick Amp Labeling Kit (Agilent, Santa Clara, CA), and then hybridized to a Gene Expression Large Volume Hybridization Kit (Agilent). Hybridized genes were

scanned by Gene chip scanner 3000 system. Data were analyzed by GeneSpring software (Agilent Technology, Santa Clara, CA).

Flow cytometry

The medium was removed after each incubation, and cells were re-stimulated with 50 ng/ml phorbol 12-myristate 13-acetate (PMA) (Sigma) and 500 ng/ml ionomycin (Sigma) in the presence of GolgiPlug (BD bioscience) for 4 h, followed by fixation and permeabilization with a fixation/permeabilization solution (Cytofix/Cytoperm, BD Pharmingen). Cells were then stained with anti-mouse CD45.2 antibody (eBioscience), anti-mouse CD4 antibody (BioLegend), anti-mouse IFN- γ antibody (eBioscience) and anti-mouse IL-17A antibody (BioLegend) followed by FACS analysis on LSR Fortessa (BD Bioscience).

Western blot analysis

117 Differentiated Th17 cells were cultured with 10 ng/ml IL-23 for 3 days to induce
118 IL-23R, rested to return STAT3 phosphorylation to the basal level, and then
119 re-stimulated with 100 μ M db-cAMP and/or 100 ng/ml IL-23 for 30 min with indicated
120 compounds for indicated time. Total cell lysates were prepared with RIPA buffer
121 (Sigma) containing a phosphatase inhibitor cocktail (PhosSTOP, Roche, Basel,
122 Switzerland) and a proteinase inhibitor cocktail (Complete Protease Inhibitor Cocktail,
123 Roche). Lysates were then subjected to SDS-PAGE (sodium dodecyl
124 sulfate-poly-acrylamide gel electrophoresis) and separated proteins were transferred to a
125 PVDF membrane (Millipore, Darmstadt, Germany). After blocking with an ECL
126 Blocking Agent (GE Healthcare, Piscataway, NJ), membranes were incubated with
127 primary antibodies, followed by incubation with secondary antibodies conjugated with
128 horseradish peroxidase (GE). Signals were detected using an ECL Prime Western
129 Blotting Detection Reagent (GE) on LAS-4000 (GE). Primary antibodies used were;
130 mouse monoclonal anti- α -Tubulin antibody (clone DM1A, #T6199, Sigma), mouse
131 monoclonal anti-GAPDH antibody (clone 6C5, #AM4300, Ambion, Austin, TX), rabbit
132 monoclonal anti-STAT3 antibody (clone 79D7, #4904, Cell Signaling Technology,

Danvers, MA), rabbit monoclonal anti-phosphorylated STAT3 antibody (Y705; #9145, Cell Signaling Technology), rabbit monoclonal anti-phosphorylated STAT3 antibody (S727; #9134, Cell Signaling Technology), rabbit monoclonal anti-JAK2 antibody (clone D2E12, #3230, Cell Signaling Technology), rabbit monoclonal anti-phosphorylated JAK2 antibody (Tyr1007/1008; #3771, Cell Signaling Technology), rabbit monoclonal anti-NF- κ B p65 antibody (clone D14E12, #8242, Cell Signaling Technology), rabbit monoclonal anti-phosphorylated NF- κ B p65 antibody (S536; clone 93H1, #3033, Cell Signaling Technology), rabbit monoclonal anti-p105/p50 antibody (#3035, Cell Signaling Technology), and rabbit monoclonal anti-phosphorylated p105 antibody (S933; #4086, Cell Signaling Technology).

RNA interference

siRNA for mouse *Creb1* (5' -UUGAACAACAACUUGGUUGCUGGGC-3' (sense) or 5' -GCCCAGCAACCAAGUUGUUGUCAA-3' (antisense) and scrambled control siRNA were obtained from Invitrogen (Stealth RNAi, Carlsbad, CA). Th17 cells differentiated with TGF- β 1 (1 ng/ml) and IL-6 (20 ng/ml) were transfected with 500

149 pmol of each siRNA using an Amaxa P3 Primary Cell 4D-Nucleofector X Kit with the
150 program DN100 on a 4D-Nucleofector (Lonza, Basel, Switzerland) in 100 μ l. After
151 transfection for 4 h, cells were washed and stimulated with or without 10 ng/ml IL-23
152 for 2 days , and then incubated with or without 100 μ M db-cAMP for 1 day. Total RNA
153 was prepared and then subjected to qRT-PCR analysis.

154

155 **Histology**

156 The ear tissues from psoriasis models were removed and fixed by 4 %
157 Paraformaldehyde (PFA) for 48h at 4°C. Each ear tissues were embedded in paraffin,
158 sectioned at 5 μ m thickness, and then stained with hematoxylin-Eosin.

159

160 **Analysis of gene expression of human skin biopsies and mouse IL-23-treated ear**

161 **from microarray datasets**

162 Microarray gene expression data of human skin biopsies were retrieved from Gene
163 Expression Omnibus datasets (GSE51440 and GSE13355).^{52,53} Patients information and

skin samples have been described previously.^{52,53} In brief, two biopsies were taken from each patient - one from lesional skin of each patient (involved sample) and the other from non-lesional skin (uninvolved sample), taken at least 10 cm away from any active plaque. One biopsy was obtained from each healthy control.⁵² Microarray gene expression data of IL-23-treated ear from mice was retrieved from GSE13335.⁵³ Gene expression levels were transformed to z-score values. *P* values were calculated by nonparametric Wilcoxon-Mann-Whitney test, and correlations between expression levels of two genes were calculated by nonparametric Spearman correlation test.

Statistical Analysis

Data are shown in mean \pm SEM. Statistical comparisons among more than two groups were conducted using One-way ANOVA with Bonferoni test. Statistical comparisons between two groups were conducted using Mann-Whitney test. *P* values of 0.05 or less were considered significant.

179 **REFERENCES**

- 180 E1. Sugimoto Y, Narumiya S. Prostaglandin E receptors. *J Biol Chem*
181 2007;282:11613-11617. <https://doi.org/10.1074/jbc.R600038200>
- 182
- 183 E2. af Forselles KJ, Root J, Clarke T, Davey D, Aughton K, Dack K, et al. In vitro and
184 in vivo characterization of PF-04418948, a novel, potent and selective prostaglandin
185 EP(2) receptor antagonist. *Br J Pharmacol* 2011;164:1847-1856.
186 <https://doi.org/10.1111/j.1476-5381.2011.01495.x>

ProbeName	GeneSymbol
A_51_P193686	1700012B09Rik
A_55_P2059010	Rbp1
A_66_P116173	Il23r
A_30_P01019068	
A_55_P2424921	1300014J16Rik
A_30_P01030266	
A_30_P01021272	
A_55_P2059765	Foxf1
A_55_P2300071	4833412C15Rik
A_30_P01019394	
A_55_P2156697	Il17a
A_30_P01033385	
A_30_P01020711	
A_51_P519301	Il17f
A_30_P01023418	
A_52_P536494	Mycn
A_55_P2139942	Calca
A_51_P415306	4930563D23Rik
A_52_P374653	
A_51_P435844	Nr2e3
A_30_P01028775	
A_51_P271503	Il1r1
A_55_P1995647	Rsph4a
A_52_P593465	Arap2

ProbeName	GeneSymbol
A_55_P1962523	Zic2
A_52_P650387	Ccnjl
A_51_P146970	Dmrt2
A_51_P167527	Lum
A_51_P290387	Sval1
A_55_P2199202	Il22
A_55_P1985850	Timp1
A_55_P2089957	Myo3a
A_30_P01025708	
A_55_P1986647	Podxl2
A_30_P01032975	
A_30_P01023119	
A_55_P1962400	Il1rn
A_55_P1994942	Rorc
A_55_P1980771	Olfr536
A_55_P2052016	Crispld2
A_55_P2055597	Enpp2
A_52_P415155	Wnt6
A_55_P2238965	Cwh43
A_30_P01029390	
A_55_P1998737	9130015A21Rik
A_51_P212754	Tgfb1
A_55_P1991465	

ProbeName	GeneSymbol
A_55_P2153545	
A_55_P1955078	Igflr1
ERCC-00018_67	
A_51_P483576	
A_55_P2372228	A430104N18Rik
A_55_P2040490	
A_55_P2130104	Slc52a3
A_55_P2149791	
A_55_P2224830	D230044B12Rik
A_30_P01021682	
A_55_P2011341	
A_55_P2091496	Dppa3
A_55_P1977558	Dip2b
A_30_P01027915	
A_55_P2149500	Kifc2
A_55_P2175502	M1ap
A_51_P288916	Tmtc2
A_52_P21550	Gcnt1
A_52_P221776	Kif12
A_30_P01028727	
A_30_P01032107	
A_55_P1957865	
A_51_P211854	Selp
A_55_P2041514	4930550L24Rik
A_55_P2141093	Eya2
A_30_P01031172	
A_55_P2180176	Ms4a6b
A_55_P2004511	Cd300lf
A_55_P2049687	Efna2
A_55_P2322555	5930433N17Rik
A_30_P01026837	
A_30_P01030563	
A_66_P130030	LOC102642739
A_30_P01027181	
A_30_P01023491	
A_30_P01026677	
A_52_P16232	Gabbr1

A_30_P01022661	
A_30_P01022155	
A_30_P01021193	
A_52_P214408	Gm5148
A_51_P386670	Dse
A_55_P2156598	Gm9869
A_55_P1993483	
A_66_P101393	A530001N23Rik
A_51_P132013	Cysltr2
A_51_P407657	
A_66_P102202	Gm648
A_30_P01024802	
A_52_P244193	Cd24a
A_51_P108459	Gpr65
A_55_P2039354	
A_55_P2404823	1700095A21Rik
A_55_P2101944	
A_30_P01028859	
A_51_P124748	Tgfb3
A_30_P01018979	
A_55_P1996434	
A_55_P2129207	Tmem71
A_66_P124806	Tlr4
A_30_P01019442	
A_55_P2148873	Cat
A_30_P01031480	
A_55_P2344598	E330037I15Rik
A_30_P01019859	
A_51_P505823	Endod1
A_30_P01025027	
A_55_P2397599	B930025B16Rik
A_30_P01024408	
A_30_P01019299	
A_30_P01025650	
A_55_P2409088	BB163080
A_55_P2006808	Ntrk3
A_30_P01021307	
A_30_P01019924	

A_55_P2051270	Tctex1d1
A_30_P01027278	
A_55_P2069550	
A_55_P2113310	
A_55_P2213348	5330421C15Rik
A_30_P01019195	
A_55_P2106358	
A_30_P01026021	
A_55_P2027392	Gpr146
A_55_P2100290	Adra1b
A_55_P1976200	Lmtk3
A_51_P111612	Arrdc4
A_30_P01029744	
A_30_P01031674	
A_30_P01024836	
A_55_P2132646	
A_51_P423976	Crem
A_65_P01319	Pde4b
A_55_P2001262	
A_51_P484998	Hgf
A_55_P2178653	
A_55_P2416087	C530005A16Rik
A_55_P2040600	Exd1
A_51_P117581	Cables1
A_55_P2157033	Bace2
A_55_P2144095	
A_55_P2127303	Spsb2
A_55_P2022629	Oxct2b
A_55_P2108248	Art4
A_30_P01032657	
A_55_P1954835	Ramp3
A_55_P1973427	Tex14
A_55_P2053838	Tnfaip3
A_51_P491350	Col4a2
A_51_P438990	Olfr911-ps1
A_55_P2015887	Wwox
A_55_P2224431	6330412A17Rik
A_51_P200544	Tnip3

A_55_P2153391	Palm2
A_55_P2090374	Vmn1r68
A_66_P106388	Ms4a4c
A_55_P1958038	Klra16
A_55_P2003813	Scn3b
A_55_P2270412	C230096K16Rik
A_30_P01033617	
A_52_P799815	Tmem171
A_55_P2195172	D930021N14
A_51_P297069	Tmod1
A_30_P01019014	
A_30_P01018802	
A_55_P2148809	Zan
A_55_P2042615	
A_51_P505868	Lhfp
A_55_P2013184	Atp2b3
A_51_P186798	
A_55_P1969962	Catsper3
A_30_P01020171	
A_66_P109183	Apold1
A_55_P1996911	Rasa3
A_55_P1964896	Gm3014
A_55_P2256586	Gm11783
A_55_P2259896	A130009E19Rik
A_52_P541833	Vps37b
A_30_P01018712	
A_51_P121915	BC089597
A_30_P01028426	
A_55_P2126259	Slc35e2
A_55_P2174323	Pgpep1l
A_30_P01018806	
A_55_P1955183	Crxos
A_30_P01017739	
A_55_P1963046	Prl3c1
A_55_P2020361	Lzts3
A_51_P422685	Zmat4
A_66_P139805	Crispld1
A_55_P2181191	Btg1

A_55_P2168223	Aicda
A_55_P2381821	6430706H07Rik
A_30_P01023753	
A_55_P2120064	Gm2984
A_30_P01028595	
A_55_P2269254	5730419F03Rik
A_30_P01027938	
A_55_P2128734	Pbxip1
A_51_P106249	Cmtm2b
A_55_P2143516	
A_55_P1978416	Il12rb2
A_55_P1968763	Shc4
A_55_P2113673	Eml1
A_51_P112223	Gsta4
A_55_P2020035	
A_55_P2016064	
A_55_P2419021	9330188P03Rik
A_55_P2321453	D11Erttd726e
A_52_P1093529	Pik3r5
A_55_P1981829	Rhox8
A_51_P170959	Proz
A_30_P01031286	
A_51_P123676	Synpo
A_30_P01032395	
A_30_P01020338	
A_52_P577388	Epdr1
A_65_P10913	Tgfb2
A_30_P01031663	
A_55_P2069802	D5Erttd577e
A_52_P57317	Fam19a3
A_55_P1977628	Pappa
A_55_P2146254	Ifitm1
A_30_P01026001	
A_52_P460957	Crem
A_55_P2234084	
A_30_P01030451	
A_55_P2017826	Myb
A_30_P01030387	

A_55_P1962284	Klhl24
A_30_P01023132	
A_30_P01020686	
A_51_P234253	Sdcbp2
A_51_P229664	Cd27
A_55_P1966568	
A_30_P01020841	
A_52_P22763	Map2
A_30_P01022829	
A_30_P01025657	
A_55_P2131168	Sv2c
A_66_P117730	Hapln1
A_55_P2156638	Gpr114
A_51_P264825	Lag3
A_30_P01032069	
A_55_P2162543	
A_30_P01017626	
A_52_P190973	Vcl
A_52_P491872	D830046C22Rik
A_51_P253883	Fam49a
A_51_P480190	Spaca1
A_52_P665675	Abca1
A_51_P206405	Ptprz1
A_30_P01023097	
A_30_P01022080	
A_55_P2087963	
A_55_P1976471	
A_51_P405397	Ecm1
A_55_P1987725	Gria4
A_51_P394847	Gm11346
A_30_P01022578	
A_30_P01018048	
A_55_P2133632	Sipa1l2
A_52_P163924	Trpd52l3
A_30_P01022282	
A_55_P2058270	LOC101056056
A_55_P2149931	Arap2
A_55_P1978696	Abcb4

A_55_P1991381	Adam34
A_66_P124179	Atp6v0d2
A_55_P2134877	Eml1
A_55_P2257076	
A_55_P2069485	Ptpn13
A_66_P105801	Igf1r
A_55_P1975690	Bicc1
A_55_P2148935	
A_55_P2007646	Cryaa
A_51_P340170	Il20ra
A_55_P2134246	Serpnb9
A_30_P01024596	
A_55_P2090330	Kcnmb4
A_30_P01019779	
A_55_P2004551	Klra1
A_30_P01018759	
A_55_P2096127	
A_55_P2370250	Syn3
A_30_P01031842	
A_55_P2118866	Cmah
A_55_P1962004	
A_55_P2074144	Tmprss11e
A_30_P01019251	
A_55_P1989772	Sqrdl
A_55_P1986596	Cacna1h
A_66_P101261	Gm3367
A_55_P2143219	Rasgrp2
A_55_P1968024	
A_55_P2021398	
A_55_P1975045	Sgcg
A_30_P01031226	
A_30_P01027578	
A_55_P2087265	Ifitm1
A_55_P2030672	Gm1587
A_30_P01020408	
A_30_P01021308	
A_55_P2249379	Gm16982
A_55_P2115567	Slc26a1

A_30_P01030530	
A_30_P01029489	
A_51_P185292	4930581F22Rik
A_55_P2174847	Olfr566
A_52_P56682	Sla2
A_55_P2056533	Ntrk3
A_55_P2129407	4932431P20Rik
A_55_P1989765	
A_51_P470079	Il1r2
A_55_P2113703	Spib
A_51_P352452	Dbx2
A_55_P1960846	4933408B17Rik
A_55_P2144526	Fam65b
A_55_P2112693	Sipa1l1
A_55_P2110497	Ddc
A_30_P01033126	
A_55_P2117164	Tmem106b
A_55_P2157134	Rmnd5a
A_55_P2367255	4833421G17Rik
A_55_P2132781	Slc16a2
A_52_P68702	Frmd4b
A_30_P01022421	
A_30_P01032372	
A_51_P254646	Jdp2
A_55_P2035843	
A_55_P2165199	Cxcr6
A_55_P2172999	Ptpn13
A_55_P2105239	Defb47
A_51_P460279	Fam154a
A_55_P1989813	Gcm1
A_51_P429335	Prss16
A_55_P1964594	
A_55_P1961608	Ypel4
A_55_P1985831	
A_30_P01029085	
A_55_P2029902	Gab3
A_55_P2129469	
A_51_P414126	Rab19

A_55_P2185890	Cfh
A_55_P2011390	Tead1
A_55_P2062058	Dbnidd2
A_30_P01028164	
A_66_P109986	Cd33
A_30_P01029324	
A_55_P2150476	
A_55_P2077522	
A_51_P483280	Prnp
A_55_P2277620	Gm16523
A_30_P01022763	
A_55_P2268790	4930445G23Rik
A_55_P2258567	D930043N17Rik
A_55_P2019949	
A_52_P585124	Cxcr4
A_55_P1968789	Rasa3
A_55_P2030030	Adssl1
A_55_P2013128	
A_55_P2070494	Vmn2r60
A_55_P2038347	Acot3
A_66_P111534	5430431A17Rik
A_55_P2137049	AA467197
A_30_P01022950	
A_30_P01027369	
A_55_P2058601	LOC102636451
A_55_P2152771	Lhfpl2
A_55_P2027979	Impg2
A_55_P2156126	Sema6c
A_30_P01027097	
A_55_P2066553	Oacyl
A_52_P236398	
A_52_P203560	Fzd10
A_55_P2028837	Tspan2
A_51_P451458	Mamdc2
A_55_P2058671	Zfyve28
A_30_P01025472	
A_55_P2055189	Kcng1
A_55_P1991874	Dcpp3

A_55_P2361932	Far2os2
A_55_P2107247	Tssk5
A_30_P01031426	
A_55_P1962699	
A_55_P2050747	4930557B15Rik
A_51_P242930	Lat2
A_51_P463187	Tbc1d2b
A_30_P01021805	
A_55_P2058195	Fsip2
A_55_P2299275	Olfr22-ps1
A_52_P108850	St8sia1
A_30_P01032338	
A_52_P173442	Wscd2
A_55_P2438722	Ifnar1
A_30_P01026506	
A_30_P01021489	
A_51_P262721	0610009L18Rik
A_55_P2068515	Gm5083
A_55_P2064351	Vipr1
A_30_P01028109	
A_55_P2147487	Cyth3
A_51_P103541	Cacna1s
A_30_P01033260	
A_55_P1971009	Gzme
A_30_P01024586	
A_55_P2016114	Fasl
A_65_P18181	Runx2
A_52_P14526	Zyg11b
A_55_P2167930	Dmd
A_55_P2121595	Scml4
A_55_P2410325	2610028D06Rik
A_55_P2059657	
A_30_P01023293	
A_55_P2199118	Bend4
A_30_P01032207	
A_30_P01031205	
A_55_P1967002	
A_30_P01032896	

A_55_P2133165	Wwc1
A_55_P2118609	St6galnac1
A_52_P663526	Nmrk1
A_55_P1981719	Rreb1
A_30_P01033663	
A_52_P585907	Pla2g4f
A_55_P2180854	Mrgprg
A_30_P01027293	
A_30_P01029470	
A_55_P2107383	Vmn2r96
A_55_P2401971	AU015680
A_55_P2068248	
A_55_P2057577	Ugt1a6a
A_55_P2098175	Olfr701
A_52_P489778	Ablim1
A_55_P2278531	C920008N22Rik
A_55_P2146683	Gm4657
A_55_P2014388	Olfr54
A_55_P1960479	
A_55_P1968028	Tdgf1
A_30_P01019129	
A_51_P155323	Hc
A_55_P2192397	A330023F24Rik
A_55_P1974178	Pyhin1
A_55_P2091359	Padi2
A_51_P461067	
A_55_P2062250	Gm5524
A_30_P01022466	
A_30_P01018533	
A_52_P156452	Cmah
A_30_P01029190	
A_51_P363749	Irf6
A_51_P368823	Grb7
A_55_P1957413	Lsp1
A_52_P574668	Nt5e
A_55_P1989514	2010016I18Rik
A_30_P01027566	
A_30_P01024360	

A_30_P01029607	
A_55_P2183208	PrI2c1
A_52_P212686	Lrrk1
A_52_P162500	Dph3
A_30_P01027605	
A_51_P382152	Procr
A_65_P12530	Pde4d
A_55_P1981200	Grm1
A_55_P1988975	Ms4a4b
A_55_P2184945	Mageb16
A_30_P01031942	
A_66_P108059	Ttc39c
A_51_P484111	Matn2
A_55_P2165249	Papln
A_55_P2228953	4633402D09Rik
A_30_P01018318	
A_55_P2130660	Slitrk4
A_52_P532227	S1pr1
A_30_P01030847	
A_30_P01028877	
A_30_P01018107	
A_52_P140005	Nipal1
A_52_P622850	Hes5
A_51_P176972	Amigo2
A_30_P01029731	
A_55_P2040497	Gm11517
A_55_P2123566	Sipa1l2
A_55_P2161045	
A_52_P390127	Klrc1
A_52_P118706	9630013D21Rik
A_55_P1961761	Dcc
A_51_P416858	Myl1
A_66_P114641	Cdh18
A_30_P01030359	
A_51_P215475	Ptprb
A_51_P246773	Sesn3
A_55_P2106763	Pxdc1
A_30_P01019330	

A_51_P493987	Moxd1
A_30_P01023895	
A_30_P01019088	
A_30_P01033069	
A_55_P1994289	Gm10791
A_52_P553890	Itgb3
A_30_P01028896	
A_55_P2054409	Pira2
A_30_P01032087	
A_55_P2353079	LOC102631735
A_55_P2008093	Vmn2r91
A_55_P2023161	Gm13084
A_30_P01029619	
A_30_P01023506	
A_66_P111562	Ccnd1
A_30_P01017761	
A_30_P01031709	
A_52_P23225	Gpc3
A_30_P01027076	
A_30_P01023050	
A_55_P2117465	Gm6559
A_55_P2243828	LOC552901
A_30_P01021882	
A_52_P602669	Serpinb6d
A_55_P2106583	
A_55_P1974932	Olfr668
A_55_P1954331	Murc
A_30_P01019065	
A_55_P2113384	
A_30_P01019463	
A_55_P2135986	Ms4a4c
A_55_P2103948	
A_55_P2186605	Cacnb2
A_66_P126293	Itgb3
A_55_P2105416	Gm10319
A_51_P466731	Olfr959
A_51_P250058	Epas1
A_55_P2047356	Gm4868

A_55_P2145224	
A_55_P2272748	
A_30_P01020519	
A_55_P1973833	Astl
A_52_P836852	Txk
A_52_P71686	Atp6v0d2
A_55_P2069052	Sacs
A_30_P01025429	
A_30_P01032232	
A_55_P2283116	4930478P22Rik
A_30_P01022191	
A_55_P2099620	Hmx2
A_55_P2044922	Gpr82
A_30_P01030786	
A_51_P142896	Cd59a
A_55_P2287260	Snap23
A_55_P2048224	
A_30_P01018395	
A_52_P58359	Tlx1
A_55_P2020338	Scml4
A_30_P01031489	
A_30_P01024761	
A_55_P2131143	Dcaf17
A_66_P116314	
A_55_P1997450	
A_51_P487487	Speer4d
A_51_P149267	LOC101056131
A_55_P2108012	Fam78b
A_30_P01023920	
A_55_P2086954	
A_30_P01017444	
A_30_P01020712	
A_55_P2112005	Tff1
A_55_P1972659	Spaca1
A_55_P2032009	Cc2d2b
A_55_P2104835	P2rx4
A_55_P2330173	4930474G06Rik
A_30_P01022638	

A_55_P2248150	4930405A10Rik
A_30_P01030603	
A_51_P391445	Ifngr1
A_55_P2061273	Tbx6
A_55_P2026894	
A_55_P2155146	Klrc3
A_30_P01030998	
A_55_P2162935	Ntn1
A_30_P01032852	
A_55_P2017677	Cap2
A_30_P01018939	
A_30_P01028549	
A_55_P2020726	Gm16505
A_55_P2030903	Rsph4a
A_55_P2076744	
A_30_P01018535	
A_55_P2022504	1700019B21Rik
A_30_P01027098	
A_55_P2037962	Trpm6
A_55_P1983508	Nr4a2
A_55_P2094060	Gzma
A_55_P2032489	
A_30_P01025834	
A_52_P614207	
A_30_P01027590	
A_30_P01017808	
A_30_P01030286	
A_55_P2082319	BC094916
A_55_P2391674	4931407E12Rik
A_52_P172014	Ramp1
A_30_P01018833	
A_51_P471088	Vmn1r24
A_55_P2092492	Il18r1
A_55_P2129771	Prrt1
A_30_P01031851	
A_55_P2112772	Gabra3
A_55_P2150377	
A_30_P01019182	

A_30_P01021265	
A_55_P2101340	Ramp3
A_51_P312121	Xdh
A_55_P2024046	Slc16a5
A_55_P1990398	Vmn2r48
A_52_P500274	Ntrk3
A_30_P01024455	
A_55_P2054261	C2cd4b
A_30_P01020856	
A_55_P2056403	Speer4f
A_52_P483336	Ms4a1
A_55_P2257765	Gm7111
A_30_P01022071	
A_30_P01032178	
A_30_P01024457	
A_55_P2031403	Smim24
A_30_P01019202	
A_55_P2045928	Btnl10
A_30_P01025230	
A_30_P01022726	
A_55_P2014516	
A_55_P2164253	2010015L04Rik
A_30_P01029367	
A_51_P381618	Pla1a
A_30_P01024417	
A_30_P01022468	
A_55_P2149288	Cmya5
A_30_P01029943	
A_30_P01020320	
A_30_P01026503	
A_55_P1961567	Ypel3
A_30_P01019556	
A_30_P01029037	
A_55_P2162815	Fbxo31
A_55_P2035509	Pyhin1
A_30_P01028513	
A_52_P514407	Klra15
A_30_P01030158	

A_30_P01030095	
A_55_P2064741	Nmb
A_30_P01018793	
A_55_P2105436	Foxp1
A_51_P419246	5830416P10Rik
A_55_P1988368	Upp1
A_55_P2017169	
A_30_P01022733	
A_55_P1979950	
A_52_P161495	Bcl6
A_52_P570820	Slc22a22
A_30_P01024947	
A_51_P298066	Clnk
A_55_P2237440	Gm13999
A_30_P01028422	
A_30_P01021760	
A_30_P01023427	
A_55_P2103060	
A_55_P2121484	Smo
A_55_P2089342	Gm13547
A_52_P37091	Magi1
A_30_P01031833	
A_55_P1981830	
A_52_P648715	Triml1
A_55_P2035986	E330020D12Rik
A_55_P2081590	Olfr1320
A_55_P2108933	Dao
A_55_P1959525	Wbscr25
A_30_P01020636	
A_55_P2184434	Eomes
A_30_P01024775	
A_30_P01024742	
A_55_P2070296	D330025C20Rik
A_55_P2008016	Armc3
A_55_P2149921	
A_30_P01028539	
A_55_P2208682	B230334C09Rik
A_30_P01032937	

A_55_P1973501	Ceacam16
A_52_P197402	Tbc1d30
A_51_P247168	Wdr96
A_55_P2160416	Acox1
A_51_P202801	Abcb9
A_52_P554650	Gzmd
A_55_P2011445	Capn13
A_55_P1962209	Cxcr6
A_66_P135800	
A_51_P110341	Scgb3a1
A_52_P500979	Pcsk1
A_55_P1967591	
A_55_P2023290	1110032A03Rik
A_30_P01028087	
A_55_P2058127	Pde4dip
A_55_P1984401	Shank2
A_30_P01032347	
A_51_P456870	Foxj1
A_55_P1998892	Smox
A_51_P463765	Timp3
A_55_P2197638	1110046J04Rik
A_66_P140688	
A_66_P103231	
A_30_P01023911	
A_55_P2010396	Pde4d
A_30_P01023370	
A_51_P317640	Tgfb2
A_55_P2133943	
A_55_P2227154	A530041M06Rik
A_30_P01031611	
A_51_P419637	Dclk3
A_52_P370935	Glci1
A_55_P2063237	Dusp5
A_30_P01026901	
A_30_P01028225	
A_55_P2206269	9130403I23Rik
A_55_P1957867	Gm3161
A_30_P01018083	

A_52_P355084	Metrn1
A_51_P136294	Ms4a4b
A_55_P2035029	Mup-ps12
A_51_P285047	Cd160
A_55_P1984322	BC042761
A_66_P118299	Gm6846
A_55_P1962084	
A_55_P2084308	Nid1
A_55_P2060269	1700091H14Rik
A_55_P2004652	Klrc1
A_55_P2093614	Dennd4a
A_51_P438967	Gpnmb
A_55_P1954998	Phf1
A_55_P2122020	Klf4
A_30_P01030639	
A_51_P483013	Arel1
A_30_P01031710	
A_55_P2000439	Ptprz1
A_55_P2078123	Rora
A_55_P2066299	Gpr137b
A_66_P119376	Kctd12
A_30_P01023079	
A_66_P139530	Gm5107
A_51_P212420	Lama4
A_30_P01018590	
A_66_P136844	Dppa2
A_30_P01033579	
A_55_P2041457	
A_55_P2024150	Slco6b1
A_55_P2417434	E130215H24Rik
A_30_P01022989	
A_55_P2054362	Cx3cr1
A_55_P2290914	Trav3n-3
A_55_P1957209	4932443I19Rik
A_55_P2361652	C230085N15Rik
A_51_P372743	Frmpd3
A_55_P2141754	Tcl1b2
A_55_P2060991	BC005764

A_55_P2007339	Gm16489
A_55_P2288670	E130118H10Rik
A_55_P2033105	Cdcp1
A_30_P01032241	
A_55_P2402258	4930432J09Rik
A_55_P2081488	Pglyrp1
A_55_P2332731	
A_55_P2064928	Nebi
A_51_P497317	Tcp11l2
A_30_P01025765	
A_55_P2202524	9330162012Rik
A_66_P122719	
A_55_P2079269	
A_52_P566396	Rnf122
A_51_P120830	Mmp10
A_55_P2144556	Flrt3
A_30_P01022541	
A_30_P01017796	
A_30_P01020946	
A_55_P1961429	Vmn1r131
A_66_P117484	Gm14317
A_51_P142153	Filip1l
A_51_P281778	Igsf23
A_51_P505617	Il18r1
A_55_P1966432	Gstm1
A_55_P2076196	Mup17
A_30_P01018690	
A_30_P01029325	
A_55_P2006852	Dgkb
A_51_P420128	Fam219aos
A_52_P456134	Dgat1
A_52_P660477	Spo11
A_55_P2036086	
A_51_P132170	Ccdc141
A_51_P448127	Khdc3
A_30_P01025555	
A_55_P2175925	
A_55_P2026779	1700091H14Rik

A_51_P295967	Proc
A_30_P01019196	
A_30_P01022138	
A_55_P2223851	
A_51_P469688	Syt17
A_30_P01024849	
A_30_P01023056	
A_30_P01019238	
A_51_P259975	Aspa
A_55_P1987291	4833424O15Rik
A_30_P01018325	
A_55_P2138739	
A_30_P01018311	
A_55_P2161219	Thsd7a
A_55_P2022519	Tmem108
A_30_P01024167	
A_51_P196695	Il7r
A_55_P2084739	Gcnt1
A_30_P01029056	
A_55_P1971010	Gzme
A_52_P423247	Pde4b
A_30_P01032466	
A_55_P1960999	Pigr
A_30_P01027371	
A_55_P2057528	Arl4d
A_52_P243391	Sema4f
A_55_P2212006	2410087M07Rik
A_51_P290921	Syt12
A_51_P499838	Bst1
A_55_P2062449	Pde6a
A_55_P2058297	Sgip1
A_55_P2156140	Tcerg1l
A_55_P2307496	
A_51_P448147	Gimap7
A_55_P2214487	E330013P08Rik
A_51_P205385	Uox
A_51_P149714	Ms4a6d
A_55_P2279498	4933438K21Rik

A_55_P2070331	5830416P10Rik
A_51_P490795	Mxd1
A_52_P244702	Tcf7
A_52_P117408	Tg
A_55_P2113498	Klrd1
A_55_P2419299	1110006E14Rik
A_55_P2080476	
A_51_P330452	Olfr1414
A_52_P42194	Svil
A_51_P493117	Slc16a9
A_30_P01027848	
A_52_P605517	Phactr1
A_55_P2007831	Ccdc172
A_55_P2272979	Tdrd5
A_55_P2138291	Arl14epl
A_55_P1970915	Kdm4d
A_51_P290556	Nyap2
A_55_P2361437	Pitpnc1
A_55_P2067362	Dpep2
A_55_P2083297	E230025N22Rik
A_30_P01032949	
A_55_P2003266	Fam107a
A_55_P1965015	6530402F18Rik
A_52_P629895	Adh1
A_55_P2068947	Vmn1r203
A_55_P1953630	E330016L19Rik
A_55_P2200319	A630014C17Rik
A_30_P01028171	
A_55_P2063126	Inadl
A_55_P2221647	Al605517
A_55_P2167788	Gm6093
A_52_P279425	Cd96
A_55_P2004906	4930407I19Rik
A_55_P2079552	4930524N10Rik
A_55_P2237360	1700008O03Rik
A_30_P01031384	
A_55_P1968433	Agpat9
A_55_P2006265	Olfr1446

A_30_P01023045	
A_55_P2075469	Baalc
A_55_P2156806	Pom121l2
A_55_P2180744	Clstn3
A_30_P01031856	
A_55_P2004536	Klra4
A_30_P01028043	
A_52_P412506	Mup5
A_55_P2012171	Spata6
A_55_P2019699	Samhd1
A_30_P01032986	
A_51_P156158	Phlpp1
A_55_P2119377	Klri2
A_55_P2154107	Gcm1
A_55_P2004442	Nr2e1
A_30_P01028500	
A_55_P2228297	
A_66_P125770	Gm3942
A_55_P2129261	Arhgap36
A_51_P457196	Sfrp4
A_52_P649210	
A_55_P2010152	Sell
A_55_P2206461	A930006K02Rik
A_55_P1960197	P2ry14
A_55_P2057777	Fgfr1
A_30_P01023824	
A_55_P2169923	Cacna1s
A_55_P2062802	
A_55_P2223282	B130019D13Rik
A_30_P01021320	
A_30_P01027981	
A_52_P517984	Usp50
A_66_P128761	Pydc3
A_30_P01025525	
A_55_P2102838	
A_55_P2119548	Gm13298
A_55_P2136752	Ermn
A_55_P2008926	Slc17a3

A_51_P185763	Slc46a2
A_55_P2170836	Gm14149
A_55_P2046149	Kremen1
A_55_P1957871	
A_55_P2013356	Renbp
A_55_P2058028	
A_55_P2051622	Gm8479
A_55_P1971938	Atp2b2
A_66_P105596	
A_51_P473498	Gpr171
A_55_P2011491	
A_30_P01023578	
A_30_P01032461	
A_55_P2060376	Gzmg
A_51_P464900	Gabbr1
A_30_P01021842	
A_51_P446131	Gipc2
A_55_P2072000	Pcmt1d1
A_55_P2109877	Gm4718
A_55_P2174743	Akap7
A_51_P357914	Pdc
A_55_P2022773	Glcc1
A_55_P2056674	Lrrk1
A_55_P2283551	2810455B08Rik
A_55_P2025483	Rfesd
A_55_P2246344	5830407E08Rik
A_55_P1977855	Sall3
A_55_P2100197	
A_30_P01021190	
A_55_P2077501	
A_30_P01032661	
A_55_P2019838	LOC102635555
A_55_P2025937	
A_30_P01020571	
A_52_P88722	Senp7
A_55_P1964138	Pign
A_55_P2368680	Gm10741
A_51_P303180	Fam114a1

A_55_P2130627	Smim24
A_51_P187602	Serpinb5
A_52_P517098	Il18rap
A_55_P2106280	Gm10684
A_51_P456465	Cldn10
A_52_P72965	Unc80
A_65_P14951	Cblb
A_55_P2077515	
A_52_P730743	
A_55_P1989738	4930426L09Rik
A_52_P478444	
A_55_P2074656	Padi2
A_55_P2425761	C530043K16Rik
A_55_P2181963	Gm8369
A_51_P276479	4930486L24Rik
A_55_P1978316	Adamts14
A_52_P232813	Cxcl3
A_30_P01029179	
A_55_P1988658	4930558N11Rik
A_30_P01030381	
A_30_P01031149	
A_51_P413740	Ftcd
A_55_P2269289	Dgkg
A_30_P01032026	
A_55_P2378486	Kcnma1
A_30_P01027555	
A_55_P2050988	
A_51_P326229	Ddx25
A_55_P2393734	4933421H12Rik
A_55_P2008835	
A_55_P2111380	Ctnnd2
A_30_P01021778	
A_30_P01026400	
A_55_P2048448	Klra23
A_30_P01024983	
A_55_P2005470	Mfap4
A_30_P01031285	
A_30_P01024637	

A_30_P01030229	
A_55_P2062573	C1qtnf3
A_55_P2013043	Serpnb6b
A_55_P2009449	Pnma2
A_55_P2132651	Wisp1
A_55_P2021585	Tff1
A_30_P01032380	
A_30_P01023510	
A_30_P01031079	
A_30_P01020553	
A_55_P1969698	5430402E10Rik
A_55_P2335768	4831407H17Rik
A_30_P01024729	
A_51_P422540	Paqr8
A_51_P440460	Hip1r
A_30_P01017646	
A_30_P01022515	
A_30_P01021571	
A_30_P01022305	
A_55_P2109633	Tcf7
A_55_P2083629	Tle2
A_30_P01032048	
A_30_P01029579	
A_55_P2216822	4930551O13Rik
A_55_P2386256	D130062J10Rik
A_55_P2454784	lfnar1
A_51_P144264	Klf2
A_55_P2135526	Gzmc
A_55_P2315921	5330431K02Rik
A_55_P2021981	Ctsw
A_30_P01018708	
A_52_P123655	A630023P12Rik
A_55_P2364516	AA060545
A_30_P01029394	
A_55_P2178800	Ugt1a10
A_30_P01027083	
A_55_P1982747	Slc8a3
A_55_P2088375	Tnnt3

A_55_P1954555	LOC102637894
A_52_P64687	Camk2n1
A_55_P2036007	Rai2
A_51_P163953	Nsg2
A_30_P01023919	
A_30_P01018847	
A_30_P01019879	
A_30_P01032204	
A_30_P01018751	
A_52_P667477	Fyco1
A_55_P2397400	
A_55_P2373852	2310058N22Rik
A_52_P363216	Gcnt2
A_55_P2027077	Shc2
A_30_P01027939	
A_55_P2087205	9530077C14Rik
A_55_P2316682	A830011I04
A_55_P2281818	LOC433347
A_30_P01019369	
A_51_P116906	Rapgef3
A_51_P489138	Sptb
A_30_P01023736	
A_52_P381430	Tbc1d4
A_55_P2374337	A130071D04Rik
A_55_P2126269	Nmb
A_52_P681310	Plaur
A_51_P304397	Cpm
A_55_P2079560	Lilra6
A_55_P2096310	
A_51_P420577	Olfr983
A_55_P1997126	Ctse
A_55_P2000628	Dusp7
A_55_P2259456	4933425B07Rik
A_55_P2079579	Pira7
A_30_P01031257	
A_55_P2185068	Gm3002
A_30_P01031240	
A_51_P394394	Tspan2

A_55_P2255944	9130002K18Rik
A_55_P2040245	Piezo2
A_52_P612137	Runx1t1
A_55_P2025248	Mxd1
A_30_P01018930	
A_55_P2232057	AU022793
A_51_P220343	Wisp1
A_55_P1982227	
A_30_P01024650	
A_30_P01025307	
A_55_P2028847	
A_55_P2173039	
A_30_P01028959	
A_66_P102878	Olfr726
A_55_P1965564	Gm15085
A_55_P2156515	
A_55_P2161347	Acmsd
A_55_P2039646	
A_30_P01023052	
A_55_P2132888	Sdcbp
A_30_P01024653	
A_30_P01018360	
A_30_P01021347	
A_51_P183051	Upb1
A_55_P2150737	
A_55_P2107542	Pde4b
A_51_P245989	Ccr2
A_55_P2193424	6720420G18Rik
A_55_P2134616	Med12l
A_55_P2150697	Cpne4
A_55_P2373987	A730009E18Rik
A_30_P01019473	
A_52_P494622	Nr4a2
A_55_P2320263	Caln1
A_55_P2145465	D5Ert577e
A_51_P158545	Ankk1
A_66_P104815	Ecm1
A_52_P476731	Fam110c

A_30_P01031678	
A_51_P240693	Tecpr1
A_30_P01025404	
A_30_P01031292	
A_30_P01030435	
A_51_P104418	Dusp10
A_55_P2329660	Catsperd
A_55_P2007964	Cx3cr1
A_55_P2006677	Gm7969
A_55_P2218334	9430011C21Rik
A_52_P621588	Ifnlr1
A_55_P2109382	Adora2a
A_30_P01030963	

ProbeName	GeneSymbol
A_51_P124254	Col4a1
A_30_P01028640	
A_55_P2179074	Ciita
A_55_P2143025	Sema3c
A_30_P01033650	
A_55_P1983523	Cd300ld
A_30_P01031906	
A_30_P01029988	
ERCC-00138_246	
A_30_P01021841	
A_55_P2126072	
A_55_P2350553	4933405D12Rik
A_55_P2079064	Ppnr
A_30_P01033285	
A_51_P474459	Socs3
A_30_P01023629	
A_30_P01026288	
A_30_P01020935	
A_30_P01019221	
A_30_P01029761	
A_55_P2172058	Trim55
A_30_P01030183	
A_30_P01025039	
A_51_P212782	Il1b
A_30_P01027263	
A_30_P01025011	
A_30_P01023923	
A_30_P01020331	
A_30_P01023521	
A_30_P01023091	
A_30_P01021631	
A_55_P2171303	Bin3
A_30_P01029928	
A_30_P01017753	
A_30_P01027585	
A_52_P430304	Fam186b
A_30_P01026777	
A_55_P2325568	1700060J05Rik
A_30_P01033068	

A_30_P01022182	
A_55_P2052913	
A_55_P1960411	Lrrc23
A_30_P01019696	
A_30_P01020060	
A_30_P01020888	
A_55_P2091330	Olfr871
A_55_P2272830	
A_55_P2376363	4930432F04Rik
A_55_P2346859	C630007K24Rik
A_55_P2014034	Gm7285
A_52_P137765	Lmna
A_55_P2164265	Prl7c1
A_30_P01030364	
A_30_P01027780	
A_51_P127738	Scn2a1
A_55_P2383523	1700123I01Rik
A_30_P01021373	
A_51_P140803	Slco1b2
A_51_P160544	Efemp2
A_55_P1970876	Olfr389
A_30_P01021518	
A_30_P01020765	
A_66_P114333	Tlr12
A_30_P01028455	
A_30_P01032259	
A_30_P01021422	
A_55_P2085727	Stk38
A_30_P01021468	
A_55_P2419483	4732460I02Rik
A_55_P1958039	Klra16
A_30_P01018127	
A_55_P2397504	D130046C19Rik
A_55_P2198648	5830420C07Rik
A_55_P2018106	Gm14085
A_55_P1957866	Gm3161
A_55_P2199737	Dlgap2
A_52_P410685	Krt7
A_30_P01018893	
A_55_P2088178	Cbln2

A_30_P01026963	
A_55_P2026420	Pou6f1
A_55_P2066878	Kcnj13
A_55_P2128582	
A_55_P2092717	Trim43b
A_30_P01023033	
A_55_P2056876	Akap13
A_51_P475342	Chrnbl
A_55_P2249556	A630081D01Rik
A_30_P01025016	
A_30_P01031844	
A_55_P1997421	Gm5416
A_30_P01022699	
A_30_P01025043	
A_30_P01028130	
A_55_P1957277	Obfc1
A_30_P01020539	
A_55_P2403874	D230014I24Rik
A_55_P1980868	Gm6313
A_30_P01023317	
A_30_P01022249	
A_30_P01026161	
A_55_P1998827	Prim2
A_55_P2019533	Zscan4c
A_30_P01025818	
A_30_P01020406	
A_55_P2181029	
A_55_P2025038	Cpe
A_30_P01031340	
A_30_P01022117	
A_55_P2162364	Cacfd1
A_55_P2046328	Gm6225
A_66_P110091	Gm12371
A_55_P2112982	Tcl1b4
A_30_P01028297	
A_55_P2043932	Tmem8b
A_55_P2151638	Klra15
A_55_P2123123	Zic5
A_52_P401484	Inha
A_52_P434055	Birc3

A_55_P1979575	Shroom2
A_55_P2200029	Dleu2
A_30_P01032387	
A_55_P2119633	Gnal
A_66_P131931	Aurkc
A_30_P01024656	
A_52_P497021	Spred3
A_55_P2165224	4930467J12Rik
A_30_P01026625	
A_55_P2171196	
A_30_P01027893	
A_51_P260265	Hoxd4
A_52_P616332	Atp10d
A_55_P2143081	
A_30_P01027753	
A_66_P137660	Fam166b
A_55_P2148418	Vmn2r82
A_52_P51078	Ctsh
A_55_P1994290	Gm10791
A_55_P2061064	Ggt5
A_55_P2115260	
A_66_P136569	
A_30_P01028376	
A_52_P964651	Fam65c
A_30_P01026211	
A_55_P2110171	
A_55_P2108086	
A_55_P2064386	Gm4098
A_30_P01026654	
A_30_P01030450	
A_30_P01027602	
A_55_P1999756	
A_55_P2248320	C430003N24Rik
A_55_P1962289	
A_30_P01031102	
A_52_P464228	1700009N14Rik
A_66_P119841	Ddx17
A_55_P2329313	C530014P21Rik
A_30_P01019408	
A_30_P01030630	

A_65_P03728	St3gal1
A_55_P2212498	C030005K06Rik
A_55_P2162910	Rtn1
A_30_P01028299	
A_30_P01028569	
A_55_P1968895	Prph
A_55_P2409336	4932429P19Rik
A_55_P2049262	Gm16525
A_30_P01028919	
A_30_P01032993	
A_55_P1955427	Cux1
A_55_P1958329	
A_30_P01018695	
A_30_P01018313	
A_55_P1988388	Gm2347
A_55_P1957245	
A_55_P2317341	
A_51_P477682	Prss12
A_55_P2090484	Pde10a
A_55_P2250424	9530029O12Rik
A_30_P01028796	
A_30_P01030240	
A_55_P2212458	LOC548102
A_55_P2071716	Klre1
A_51_P141926	Fxyd4
A_55_P2234361	Rnf150
A_55_P2011700	Gm3364
A_55_P2086323	BC051408
A_55_P2052076	Obox3
A_51_P260740	Pcdh7
A_52_P146403	Arhgef38
A_55_P2190152	4921509J17Rik
A_51_P454873	Npy
A_55_P2048912	Themis2
A_66_P134109	Gm3331
A_30_P01027969	
A_51_P195958	Phlda1
A_65_P10029	Prdm2
A_55_P2065140	A930003A15Rik
A_52_P356093	B3galt2

A_55_P2022347	
A_51_P444447	Cebpd
A_55_P2260052	Gm17753
A_55_P2050513	Pgbd1
A_55_P2007470	Pdgfa
A_55_P2076064	Dnah8
A_55_P2008297	Cd300a
A_51_P320357	Grin2b
A_55_P2003199	Setd1b
A_55_P1964672	Krt28
A_55_P2119927	
A_55_P2127977	
A_66_P137556	Tle1
A_55_P2293351	Slc2a4rg-ps
A_30_P01017519	
A_30_P01023472	
A_30_P01033074	
A_55_P2070105	
A_30_P01019307	
A_30_P01029310	
A_30_P01022226	
A_30_P01029091	
A_66_P116860	5031434O11Rik
A_30_P01020677	
A_55_P1966155	Wfdc6b
A_30_P01023532	
A_30_P01019630	
A_30_P01028922	
A_30_P01032113	
A_51_P497741	Wdr95
A_55_P1991500	Obfc1
A_55_P2346736	A430105D02Rik
A_52_P594302	Lrba
A_30_P01019783	
A_55_P2130695	Armc4
A_30_P01024134	
A_30_P01018888	
A_30_P01028942	
A_52_P458647	
A_55_P2025765	Adam8

A_52_P266132	Fgl2
A_30_P01020996	
A_55_P1977938	Fcgbp
A_55_P2082519	Olfr883
A_30_P01032602	
A_55_P1958951	
A_30_P01021398	
A_30_P01024624	
A_55_P2075065	Gm10471
A_30_P01032240	
A_55_P2311208	C130045F17Rik
A_30_P01018774	
A_52_P667287	Cers6
A_55_P2180481	1810020O05Rik
A_30_P01022128	
A_55_P2213968	4933416M07Rik
A_30_P01019634	
A_52_P508317	Erlec1
A_55_P1977473	Dab2
A_30_P01030677	
A_55_P2051094	Rorc
A_55_P2112270	Gm6556
A_55_P2065506	
A_55_P2105403	Nrxn3
A_55_P2040873	Gm867
A_30_P01026907	
A_55_P2006625	
A_52_P468068	Tchh
A_55_P1968103	Pla2g2c
A_55_P2073905	
A_52_P655136	Nlrc4
A_66_P136801	Peg13
A_55_P2330560	1700120E14Rik
A_30_P01030273	
A_55_P2123045	Olfr453
A_55_P2066116	Bcl3
A_55_P2019362	Deptor
A_30_P01022204	
A_30_P01022054	
A_52_P463235	Ankrd33b

A_30_P01019135	
A_30_P01021718	
A_55_P2259889	LOC102632493
A_30_P01024278	
A_55_P2057941	1700049G17Rik
A_55_P2220342	C230094B09Rik
A_55_P2307578	E530011L22Rik
A_55_P1957459	Lilrb4
A_30_P01030411	
A_30_P01017989	
A_55_P2014427	Il17re
A_55_P1974622	
A_55_P2077218	Spef1
A_30_P01028994	
A_30_P01030141	
A_30_P01025370	
A_55_P1964902	Gm3014
A_55_P2249849	Sema6a
A_55_P1966470	
A_51_P184484	Mmp13
A_66_P127412	
A_55_P2126870	Nsf
A_55_P2077884	Kat6b
A_55_P2014987	Gatsl2
A_30_P01019902	
A_30_P01033039	
A_55_P2153496	Ppp2r3d
A_55_P2005984	Wfdc15b
A_30_P01031894	
A_55_P2218483	A730093L10Rik
A_30_P01033525	
A_30_P01027071	
A_51_P150745	Olfr1044
A_30_P01032097	
A_55_P2004179	Col2a1
A_55_P2053236	
A_55_P2420983	6330575P09Rik
A_30_P01026306	
A_55_P2054410	
A_55_P2110615	Slc37a2

A_55_P1958160	Sgce
A_55_P1952274	C030016D13Rik
A_30_P01018042	
A_55_P1995195	Fosl2
A_55_P2158181	Olfr39
A_55_P1967158	Gm5486
A_55_P2006479	3300002I08Rik
A_30_P01029853	
A_30_P01025615	
A_55_P2023607	Ikzf1
A_51_P486188	Pabpc2
A_55_P2099466	
A_55_P2186634	Pdik1l
A_52_P396312	Cdh17
A_55_P1979904	Mup1
A_55_P2020217	Smok4a
A_30_P01032219	
A_55_P2093874	Pbxip1
A_55_P2185332	Il17rc
A_55_P1999102	Pi16
A_30_P01025737	
A_30_P01019416	
A_30_P01022612	
A_52_P259508	Prkcq
A_30_P01022038	
A_55_P2242089	Cog5
A_55_P2014352	
A_51_P305061	0610039K10Rik
A_55_P2041240	Nav2
A_55_P2040170	Pmp22
A_30_P01030554	
A_55_P2019620	Vmn2r94
A_30_P01022174	
A_55_P2408848	Lgr4
A_30_P01020577	
A_30_P01017726	
A_55_P2102464	
A_55_P1962781	
A_30_P01017817	
A_66_P114229	Zic5

A_30_P01024104	
A_30_P01021723	
A_30_P01027051	
A_55_P2082684	Krt12
A_55_P1966608	
A_55_P2266679	Mettl21c
A_55_P2065429	
A_55_P2364315	D030002E05Rik
A_30_P01021199	
A_30_P01030589	
A_30_P01019520	
A_51_P424641	Sirt4
A_30_P01032902	
A_30_P01024889	
A_55_P2137867	Mpp3
A_55_P2177712	LOC102638515
A_55_P2105180	Bhmt
A_30_P01029327	
A_30_P01032254	
A_51_P284426	Cstad
A_55_P2043782	Trpm1
A_30_P01024577	
A_55_P2325758	A430054B03
A_30_P01031901	
A_30_P01020426	
A_30_P01023251	
A_55_P2333126	Mgea5
A_55_P1977451	
A_55_P2065726	Snx29
A_30_P01021610	
A_55_P1966770	
A_55_P2125588	Pdgfa
A_55_P2025775	
A_55_P2381921	4930406D14Rik
A_30_P01022472	
A_55_P2045158	Mrxip1
A_30_P01032514	
A_55_P1959753	Top3b
A_30_P01018341	
A_55_P2294109	1700047E10Rik

A_30_P01025798	
A_30_P01023172	
A_55_P1989812	
A_30_P01023118	
A_30_P01023385	
A_30_P01030643	
A_55_P2006008	Serpinb1a
A_52_P398989	Cytip
A_30_P01028292	
A_30_P01018167	
A_30_P01026636	
A_30_P01021342	
A_30_P01023902	
A_30_P01025624	
A_55_P2259500	D130012P04Rik
A_30_P01031220	
A_55_P2246014	C130051F05Rik
A_51_P121607	4930546C10Rik
A_30_P01026456	
A_30_P01018358	
A_55_P2086455	Wscd2
A_30_P01028737	
A_55_P2085015	Vsx1
A_51_P382849	Emb
A_55_P2139464	
A_55_P1998781	Zcchc16
A_55_P1997544	
A_55_P2075569	Smim23
A_30_P01026305	
A_55_P2122633	Airn
A_55_P1966721	9930105H17Rik
A_30_P01025549	
A_55_P2004016	Crispld2
A_55_P1958971	Gm5538
A_55_P2115062	Cdc20b
A_30_P01032903	
A_52_P38964	Sap25
A_30_P01020315	
A_66_P125948	Cass4
A_55_P2302290	E230012P03

A_55_P2276224	9330175E14Rik
A_55_P2132014	Tekt2
A_30_P01019457	
A_55_P2098941	1700110I01Rik
A_30_P01026925	
A_30_P01023697	
A_51_P238722	Cd93
A_55_P2145626	Krt82
A_51_P234692	Neat1
A_30_P01018776	
A_55_P2127854	
A_30_P01031732	
A_30_P01033290	
A_51_P411917	Gata6
A_30_P01020135	
A_55_P2141729	Snrnp27
A_55_P2261772	Lzts1
A_55_P1965313	Mctp2
A_55_P1992849	Adrb3
A_51_P303089	Ttc28
A_52_P127925	Tfec
A_30_P01025362	
A_55_P2144090	
A_30_P01026215	
A_55_P2049261	
A_30_P01033532	
A_51_P262111	Foxf1
A_52_P424231	Adamts16
A_55_P2057806	Clrn2
A_55_P2077618	Csgalnact1
A_55_P2146655	LOC102638893
A_30_P01019695	
A_55_P1972517	Gm11544
A_52_P166694	Vamp1
A_55_P2413964	1700048M11Rik
A_55_P2175976	Vmn2r35
A_55_P2147712	Ctla4
A_51_P188981	
A_55_P2168254	LOC102635322
A_55_P2171785	Dnmt3aos

A_55_P2007467	
A_55_P1958172	Ms4a5
A_55_P2211164	5330406M23Rik
A_55_P1978258	
A_55_P2040838	Gm14548
A_55_P1968355	Tle1
A_55_P1971579	Acox3
A_55_P2041584	F420015M19Rik
A_55_P2408415	Kcnq1ot1
A_30_P01029720	
A_55_P2021572	C87414
A_55_P2367415	A630026N12Rik
A_51_P477736	4932415M13Rik
A_55_P2119882	
A_30_P01028837	
A_55_P1958758	Olfr965
A_55_P2058831	Fancc
A_30_P01024460	
A_55_P2360501	
A_55_P1981366	Lamc2
A_66_P136132	
A_55_P2200618	LOC552902
A_30_P01032009	
A_55_P2012779	Rnf167
A_55_P1959348	Olfr598
A_55_P2151685	Pira11
A_55_P2180249	Mtch2
A_55_P2059035	Gm13403
A_30_P01026880	
A_51_P158814	Marveld1
A_66_P109731	
A_55_P2007981	Fmn2
A_55_P2096515	Gm10706
A_30_P01018987	
A_55_P2208260	D230019N24Rik
A_55_P1971287	Olfr1436
A_30_P01021544	
A_55_P2275402	9330177L23Rik
A_55_P2063736	Gp49a
A_30_P01022217	

A_66_P107703	Smox
A_51_P173692	Lingo4
A_55_P2172852	Ptplad2
A_30_P01023847	
A_30_P01021174	
A_55_P1954985	
A_30_P01027176	
A_55_P2156800	Ptx4
A_55_P1991994	Ptprq
A_55_P2084686	Pdik1l
A_30_P01027527	
A_30_P01021556	
A_55_P1970144	Slc16a14
A_30_P01028537	
A_30_P01019085	
A_51_P245090	Aqp3
A_55_P1988108	Mrc1
A_30_P01024483	
A_30_P01033047	
A_30_P01022552	

ProbeName	GeneSymbol
A_66_P121012	Gm6602
A_55_P2030354	
A_55_P2094040	
A_55_P2106043	Bsx
A_30_P01030933	
A_51_P111962	Bean1
A_30_P01032018	
A_55_P2130249	Sh3gl2
A_30_P01023897	

ProbeName	GeneSymbol
A_55_P2057846	Olfr943

ProbeName	GeneSymbol
A_51_P414243	Pomgnt2
A_55_P1985764	
A_51_P436201	Gart
A_51_P165934	Phb
A_55_P1964648	Btla
A_55_P2117146	Pa2g4
A_55_P2092310	Phgdh
A_51_P313761	Shmt2
A_51_P367070	Il9
A_55_P1992838	Socs2
A_55_P1986306	Ltv1
A_55_P1971774	Fsip1
A_55_P2046877	Foxq1
A_30_P01024631	
A_51_P171075	Csf2
A_52_P518997	Epha2
A_52_P63343	Ciart
A_51_P510891	Afp
A_51_P514085	Mx2
A_52_P547662	P2ry1
A_51_P208922	Stc2
A_51_P351015	Lta
A_51_P254234	Chchd4
A_51_P237865	Il4
A_55_P2067518	Slc13a3
A_55_P2171158	Hmgn1
A_51_P384318	C1ra
A_55_P2026818	Slc4a7
A_55_P2046262	Phgdh
A_51_P502152	Slc19a1
A_55_P1960157	Bcat1
A_55_P2115127	Mphosph10
A_51_P110301	C3
A_55_P2158498	Btk
A_51_P377376	Gnl3
A_55_P2175767	Rangrf
A_51_P430766	Il10
A_55_P1967443	
A_51_P107362	Socs2

A_52_P105537	Nov
A_66_P106098	Kif3a
A_66_P138319	Acox2
A_55_P2128229	
A_55_P2039027	Speer6-ps1
A_51_P115005	Edn1
A_30_P01023338	
A_55_P1973838	Slc6a9
A_55_P2100968	Dnah7b
A_55_P2053459	Timd2
A_55_P1985554	B4galt4
A_55_P2038358	Acot1
A_55_P2143042	Gm8096
A_51_P207988	Ptger4
A_52_P627816	Tgm1
A_55_P2085425	Ophn1
A_55_P1959923	Cth
A_55_P2127587	Smcr8
A_55_P2287611	4930519N06Rik
A_55_P1959521	Etv4
A_55_P2009708	
A_55_P1984655	Smtnl2
A_66_P132249	Akr1c13
A_55_P2186928	
A_51_P451957	Cpne6
A_55_P2009673	Ppp1r14d
A_55_P2008599	Pcx
A_51_P106527	Fam195a
A_55_P2096867	Gap43
A_51_P317214	Hpdl
A_55_P2085142	Spp1
A_55_P1960916	Egln3
A_51_P156434	Slc25a33
A_55_P2074736	Prkar1b
A_65_P19933	Zdhhc23
A_55_P1964302	Timm8a1
A_55_P2110037	Akap7
A_51_P317443	Cd3eap
A_51_P518163	Rrp9
A_51_P242859	Akr1c12

A_55_P2002968	Coro2a
A_52_P406828	Dkc1
A_51_P291749	Pecr
A_52_P131548	Ajuba
A_55_P2066463	Enah
A_55_P1987186	Ttll9
A_55_P2021149	Cltb
A_66_P127160	Eif2b3
A_55_P2090254	Sntg2
A_55_P1953819	Btk
A_52_P20906	Twist1
A_51_P187901	Nop56
A_52_P416327	Cd226
A_52_P246703	Ak7
A_30_P01021097	
A_55_P1960023	Trmt5
A_55_P2006118	Rbp4
A_51_P183894	Fbxo15
A_55_P2019312	Car12
A_55_P1988384	Slc7a3
A_65_P08971	F3
A_51_P172251	lfrd2
A_55_P2044917	Gpr83
A_55_P1972575	Tmeff1
A_52_P651948	Fam229b
A_51_P390538	Mpeg1
A_66_P116678	Rps8
A_51_P230507	Ell2
A_51_P300709	Srm
A_55_P1973941	Slc7a5
A_55_P2062627	Tmem238
A_51_P170758	Grwd1
A_51_P358722	Lancl3
A_55_P2158873	Ppid
A_51_P401527	Rnmtl1
A_55_P2000182	Slc5a6
A_52_P196979	Trim66
A_55_P1981994	Krt17
A_55_P1966573	Gemin4
A_51_P186053	Rrp15

A_55_P2021505	Adra2a
A_66_P129111	Nasp
A_51_P466162	C1qbp
A_51_P468249	Phex
A_55_P1996354	Wdr31
A_51_P431737	Cth
A_55_P2143311	Pdxk
A_51_P238383	B4galt4
A_55_P2052563	Id1
A_55_P2023912	
A_52_P622434	Nop16
A_55_P2235931	Cacna1c
A_55_P2011220	Armcx1
A_55_P2128929	Cc2d2a
A_55_P2061084	Wfikkn2
A_66_P136228	Rai14
A_55_P2033725	Ascl2
A_52_P359965	Cpd
A_51_P114462	Ccl17
A_66_P109192	Gm2464
A_52_P400677	AW209491
A_51_P485458	Txlna
A_55_P2057035	Slmo1
A_55_P2029319	Cd70
A_55_P1969481	Hivep3
A_55_P2017636	Thbs1
A_51_P268069	Six1
A_55_P1971889	F3
A_55_P2030486	Srsf2
A_55_P2002376	Srm
A_52_P127682	Dagla
A_51_P261164	F2rl2
A_51_P410451	Tube1
A_51_P117865	Fam20c
A_51_P513992	Spag4
A_52_P452689	Atf3
A_52_P93910	Nrp2
A_51_P180974	Prkcdbp
A_55_P1956847	Nolc1
A_51_P332676	Top1mt

A_51_P346641	Armcx4
A_52_P245766	
A_52_P503663	Ffar4
A_52_P554703	Gprin3
A_55_P2163098	Akr1c18
A_52_P344290	F2r
A_52_P596755	Dnph1
A_55_P1985840	Mettl16
A_51_P164420	Eef1e1
A_55_P2063096	Txnrd3
A_55_P1978441	Kif3a
A_51_P205779	Cd5l
A_51_P431734	Fam185a
A_52_P543040	Utp14a
A_55_P2143041	Phgdh
A_55_P1992834	Socs2
A_52_P673499	Shmt1
A_55_P2186929	
A_51_P479818	Lonrf3
A_51_P335758	Chn1
A_55_P2079669	Bcat1
A_52_P387009	Egln3
A_55_P2422248	5730420D15Rik
A_52_P366047	Rpp40
A_51_P349213	Fcrl1
A_52_P459048	2900011O08Rik
A_52_P636050	Gpatch4
A_55_P2024669	Myo6
A_52_P58949	Calcr1
A_52_P97699	D430019H16Rik
A_51_P164939	Tmem150a
A_55_P2016249	Ppid
A_55_P1995497	Atad3a
A_51_P134812	Chac1
A_55_P2201822	Naf1
A_52_P191633	Fam71b
A_51_P164203	Nme4
A_51_P296487	Lss
A_55_P1997390	Cpd
A_55_P1952638	Fgd6

A_55_P2009213	Pde7a
A_55_P1990261	Chchd6
A_55_P1959748	Asns
A_30_P01031178	
A_51_P401987	Tmem37
A_51_P104392	Rpp25
A_52_P574214	Rrp1b
A_55_P2085574	Nfix
A_55_P2367803	Il2
A_51_P432199	Sap30
A_51_P449935	Ftsj3
A_51_P378789	Cxcl13
A_55_P2099790	Nefh
A_55_P2010788	Slc6a9
A_51_P125567	Mettl13
A_52_P481279	Drc1
A_51_P301636	Kazn
A_55_P2193512	Cd226
A_55_P1952744	Timm8a1
A_51_P430630	Gpr33
A_55_P2115225	Fap
A_30_P01021389	
A_51_P487690	Ifi44
A_52_P152631	Tmem17
A_55_P2005783	Ifih1
A_52_P527800	Emilin2
A_30_P01030879	
A_51_P156438	Slc25a33
A_55_P1978052	Pet112
A_52_P248403	
A_55_P2033680	
A_55_P2168823	
A_55_P2083449	Spryd7
A_55_P2042486	Dpysl3
A_55_P2080168	Dgkk
A_51_P205326	Fam198a
A_55_P1989102	Hmgn1
A_55_P2137527	Fam183b
A_55_P2116744	Xirp1
A_55_P2184370	LOC102641088

A_51_P455647	Car2
A_55_P1964093	Rangrf
A_55_P1973995	Gm6756
A_55_P2128085	LOC102641654
A_52_P853177	Angptl2
A_52_P625640	Trim9
A_55_P1958857	Nek6
A_52_P231075	Fcrls
A_51_P401907	Gm5483
A_55_P1992490	Scg2
A_55_P1958464	
A_51_P150302	Crtam
A_55_P2134800	Cinp

ProbeName	GeneSymbol
A_51_P385099	Tnf
A_55_P1967514	Dnah7a
A_55_P2103698	Isg15
A_55_P2122075	Pdcd1lg2
A_55_P2165234	2300005B03Rik
A_55_P2187141	Pdcd1lg2
A_51_P160344	Cenpv
A_55_P2012989	Slamf7
A_55_P2163138	Tm4sf5
A_52_P157880	
A_55_P2138386	Il5
A_51_P242166	Lap3
A_55_P2205858	Col6a5
A_55_P2079020	Snhg7os
A_66_P101942	Gm9706
A_55_P2179463	Tnfsf8
A_51_P496432	Acsl1
A_51_P188281	Myf5
A_51_P113178	Fam212b
A_55_P2125972	Gorasp1
A_51_P272563	Naa25
A_66_P117543	
A_55_P1964559	Smarca5-ps
A_51_P187018	Magohb
A_52_P649561	Heg1
A_51_P270426	Egr4
A_55_P2085974	Igf1
A_52_P344978	
A_51_P184728	Cnksr3
A_66_P105460	Ccdc14
A_51_P273609	Itpka
A_30_P01028030	
A_66_P128631	Cinp
A_55_P1962214	Kpna3
A_55_P2070825	Nudt5
A_55_P1993019	
A_55_P1992814	Shq1

A_55_P2130965	
A_51_P308844	Nrn1
A_52_P13897	Hook1
A_52_P639774	Gart
A_66_P128537	lsg15
A_51_P228768	Slfn3
A_51_P516085	Dntt
A_55_P2134804	Cinp
A_51_P169624	Taf3
A_55_P2180839	Il13
A_30_P01030169	
A_30_P01018645	
A_55_P2084703	Acaca
A_51_P234627	Nubpl
A_55_P2172182	Olfr1138
A_55_P2004452	Tceal8
A_55_P2128144	Il19
A_51_P200561	4930506M07Rik
A_51_P165098	Gga2
A_52_P987201	Pdzn4
A_55_P1964960	Il33
A_55_P2123037	Olfr553
A_51_P301215	Knop1
A_30_P01025391	
A_51_P232399	Acy3
A_55_P2039684	Gpr34
A_51_P347452	Htatsf1
A_51_P215530	Rnf180
A_52_P131836	Bysl
A_55_P2082806	Trib1
A_55_P2095880	Nfix

GO ACCESSION	GO Term	p-value	corrected p-value	-logP	gene
GO:0031347	regulation of defense response	2.87E-08	1.64E-04	3.78E+00	Il1r1 Il17f Foxf1 Calca Il17a Il23r
GO:0032101	regulation of response to external stimulus	1.03E-07	1.97E-04	3.71E+00	Il1r1 Il17f Foxf1 Calca Il17a Il23r
GO:0050727	regulation of inflammatory response	7.57E-08	1.97E-04	3.71E+00	Il1r1 Il17f Foxf1 Calca Il17a
GO:0080134	regulation of response to stress	1.28E-06	0.00183381	2.74E+00	Il1r1 Il17f Foxf1 Calca Il17a Il23r

GO:1900017	positive regulation of cytokine production involved in inflammatory response	7.20E-06	0.008231741	2.08E+00	Il17f Il17a
GO:1900015	regulation of cytokine production involved in inflammatory response	1.54E-05	0.014684372	1.83E+00	Il17f Il17a
GO:0006954	inflammatory response	2.83E-05	0.020216491	1.69E+00	Il17f Calca Il17a Il23r
GO:0071345	cellular response to cytokine stimulus	2.55E-05	0.020216491	1.69E+00	Il1r1 Foxf1 Il17a Il23r

GO:0031328	positive regulation of cellular biosynthetic process	6.72E-05	0.029578676	1.53E+00	Nr2e3 Il17f Mycn Foxf1 Calca Il17a
GO:0034097	response to cytokine	5.94E-05	0.029578676	1.53E+00	Il1r1 Foxf1 Il17a Il23r
GO:0045935	positive regulation of nucleobase-containing compound metabolic process	4.74E-05	0.029578676	1.53E+00	Nr2e3 Il17f Mycn Foxf1 Calca Il17a
GO:0045944 GO:0010552 GO:0045817	positive regulation of transcription from RNA polymerase II promoter	6.37E-05	0.029578676	1.53E+00	Nr2e3 Il17f Mycn Foxf1 Il17a

GO:0051173	positive regulation of nitrogen compound metabolic process	6.47E-05	0.029578676	1.53E+00	Nr2e3 Il17f Mycn Foxf1 Calca Il17a
GO:0009891	positive regulation of biosynthetic process	7.62E-05	0.03114245	1.51E+00	Nr2e3 Il17f Mycn Foxf1 Calca Il17a
GO:0033993	response to lipid	8.48E-05	0.032340873	1.49E+00	Nr2e3 Rbp1 Il17a Il23r

GO ACCESSION	GO Term	p- value	corrected p-value	-logP	gene
GO:0005615	extracellular space	8.63E- 06	0.0250513	1.6011702	Lum Tgfb1 Wnt6 Timp1 Crispld2
GO:0031012	extracellular matrix	8.10E- 06	0.0250513	1.6011702	Lum Tgfb1 Wnt6 Il1rn Timp1 Enpp2 Il22

GO ACCESSION	GO Term	p-value	corrected p-value	-logP	
GO:0044459	plasma membrane part	3.77E-10	1.73E-05	4.76E+00	<p> Cacna1s Synpo Tgfb3 Ms4a4b Cd59a Hc Il7r Selp Cd27 Ccr2 Lag3 Cd160 Sytl2 Gpnmb Gabbr1 Vmn1r24 Gpc3 Sema4f Cd24a Klrc1 Pde4b Ms4a1 Ntrk3 S1pr1 Itgb3 Slc22a22 Nt5e Cxcr4 Pla2g4f Ifnlr1 Camk2n1 Abca1 Plaur Atp6v0d2 Dcc Tdgf1 Shc4 Rasa3 Catsper3 Atp2b2 Sgcg Il12rb2 Abcb4 Grm1 Slc8a3 Shank2 Cacna1h Gria4 Ms4a4b Scn3b Klra4 Klra1 Klrc1 Ntrk3 Cx3cr1 Slc17a3 Sell Pde4d Fasf Lzts3 Slco6b1 Trpm6 Kcng1 Ntrk3 Sgip1 Vmn1r203 Tmprss11e Kcnmb4 Adra1b P2rx4 Adora2a Sipa1l1 Gabra3 Klrd1 Klri2 Smo Slc52a3 Sdcbp Wwc1 Cyth3 Klrc3 Dmd Akap7 Clstn3 Cacnb2 Catsperd Kcnma1 Igf1r Cd33 Kctd12 Atp6v0d2 Tlr4 </p>

GO:0071944	cell periphery	1.27E-08	2.92E-04	3.53E+00	<p> Cacna1s Gpr65 Arrdc4 Synpo Tgfb3 Cysltr2 Ms4a4b Cd59a Hc Amigo2 Slc46a2 Il7r Abcb9 Ptpz1 Selp Cd27 Lat2 Ccr2 Lag3 Cd160 Sytl2 Tmod1 Cpm Olfr1414 Grb7 Tspan2 Rab19 Olfr983 Paqr8 Gpnmb Khdc3 Cldn10 Gabbr1 Olfr959 Il1r2 Vmn1r24 Gpr171 Prnp Sptb Slc16a9 Bst1 Gabbr1 Ramp1 Vcl Tbc1d30 Fzd10 Gpc3 Sema4f Cd24a Magi1 Klrc1 Svl Pde4b Fam110c Ms4a1 Ntrk3 Klra15 S1pr1 Itgb3 Sla2 Slc22a22 Nt5e Cxcr4 Pla2g4f Ifnlr1 Camk2n1 Abca1 Plaur Atp6v0d2 Txk Ramp3 Igflr1 Lsp1 Klra16 P2ry14 Pigr Dcc Cxcr6 Tdgf1 Shc4 Rasa3 Catsper3 Atp2b2 Olfr668 Sgcb Il12rb2 Abcb4 Grm1 Slc8a3 Shank2 Cacna1h Gria4 Ms4a4b Vmn2r48 Rasa3 Ptpz1 Scn3b Cd300lf Klra4 Klra1 Klrc1 Olfr1446 Ntrk3 Dgkb Cx3cr1 Vmn2r91 Slc17a3 Sell Pde4d Atp2b3 Olfr54 Wwox Fasl Cap2 Samhd1 Lzts3 Slc16a5 Slco6b1 Gpr146 Tspan2 Cdcpl Trpm6 Piezo2 Gpr82 Klra23 Efna2 Cx3cr1 Kcng1 Arl4d Fgfr1 Sgip1 Pde6a Inadl Vipr1 Vmn1r203 Ptpn13 Vmn2r60 Tmprss11e Olfr1320 Nid1 Ifitm1 Kcnmb4 Olfr701 Adra1b Ramp3 P2rx4 Vmn2r96 Pde4b Art4 Adora2a Sipa1l1 Gabra3 Klrd1 Klri2 Smo Prt1 Slc52a3 Slc16a2 Sdcbp Wwc1 Ermn Rasgrp2 Ifitm1 Cyth3 Zan Cat Klrc3 Sema6c Gpr114 Thsd7a Ntn1 Cxcr6 Dmd Cacna1s Ptpn13 Akap7 Olfr566 Clstn3 Mrgprg Cfh Cacnb2 Dgkg Snap23 Caln1 Catsperd Kcnma1 Pde4b Pde4d Olfr726 Igf1r Cd33 Cdh18 Kctd12 Atp6v0d2 Tlr4 Itgb3 </p>
------------	----------------	----------	----------	----------	--

GO:0005886 GO:0005904	plasma membrane	7.45E- 08	0.0011393	2.94E+00	<p> Cacna1s Gpr65 Arrdc4 Synpo Tgfb3 Cysltr2 Ms4a4b Cd59a Hc Amigo2 Slc46a2 Il7r Abcb9 Ptprz1 Selp Cd27 Lat2 Ccr2 Lag3 Cd160 Sytl2 Cpm Olfr1414 Grb7 Tspan2 Rab19 Olfr983 Paqr8 Gpnmb Cldn10 Gabbr1 Olfr959 Il1r2 Vmn1r24 Gpr171 Prnp Sptb Slc16a9 Bst1 Gabbr1 Ramp1 Vcl Tbc1d30 Fzd10 Gpc3 Sema4f Cd24a Magi1 Klrc1 Svll Pde4b Ms4a1 Ntrk3 Klra15 S1pr1 Itgb3 Sla2 Slc22a22 Nt5e Cxcr4 Pla2g4f Ifnlr1 Camk2n1 Abca1 Plaur Atp6v0d2 Txk Ramp3 Igflr1 Lsp1 Klra16 P2ry14 Pigr Dcc Cxcr6 Tdgf1 Shc4 Rasa3 Catsper3 Atp2b2 Olfr668 Sgcg Il12rb2 Abcb4 Grm1 Slc8a3 Shank2 Cacna1h Gria4 Ms4a4b Vmn2r48 Rasa3 Ptprz1 Scn3b Cd300lf Klra4 Klra1 Klrc1 Olfr1446 Ntrk3 Dgkb Cx3cr1 Vmn2r91 Slc17a3 Sell Pde4d Atp2b3 Olfr54 Wwox Fasf Cap2 Samhd1 Lzts3 Slc16a5 Slco6b1 Gpr146 Tspan2 Cdcpl Trpm6 Piezo2 Gpr82 Klra23 Efna2 Cx3cr1 Kcng1 Ntrk3 Arl4d Fgfr1 Sgip1 Pde6a Inadl Vipr1 Vmn1r203 Ptpn13 Vmn2r60 Tmprss11e Olfr1320 Ifitm1 Kcnmb4 Olfr701 Adra1b Ramp3 P2rx4 Vmn2r96 Pde4b Art4 Adora2a Sipa1l1 Gabra3 Klrd1 Klri2 Smo Prrt1 Slc52a3 Slc16a2 Sdcbp Wwc1 Rasgrp2 Ifitm1 Cyth3 Zan Cat Klrc3 Sema6c Gpr114 Thsd7a Cxcr6 Dmd Cacna1s Ptpn13 Akap7 Olfr566 Clstn3 Mrgprg Cfhh Cacnb2 Dgkg Snap23 Caln1 Catsperd Kcnma1 Pde4b Pde4d Olfr726 Igf1r Cd33 Cdh18 Kctd12 Atp6v0d2 Tlr4 Itgb3 </p>
-----------------------	--------------------	--------------	-----------	----------	---

GO:0009897	external side of plasma membrane	2.16E-07	0.0024761	2.61E+00	Cd59a Il7r Selp Cd27 Ccr2 Lag3 Cd24a Klrc1 Ms4a1 S1pr1 Itgb3 Cxcr4 Abca1 Il12rb2 Klra4 Klra1 Klrc1 Sell FasI Klrd1 Klrc3 Kcnma1 Cd33 Tlr4 Itgb3
GO:1902531 GO:0010627	regulation of intracellular signal transduction	1.46E-06	0.0134296	1.87E+00	Dusp10 Rapgef3 Tgfb3 Cysltr2 Phlpp1 Tnip3 Selp Cd27 Sesn3 Xdh Tgfb2 Il20ra Ecm1 Sfrp4 Prnp Hgf Il18r1 Bcl6 Fzd10 Cd24a Gcnt2 Fam110c Ntrk3 Itgb3 Sla2 Hes5 Abca1 Dcc Tdgf1 Agpat9 Rasa3 Grm1 Rasa3 Dusp7 Ntrk3 Pde4d Tnfaip3 Ntrk3 Fgfr1 C1qtnf3 Dusp5 Rora Il18r1 Adra1b P2rx4 Sipa1l1 Klf4 Sipa1l2 Wwc1 Sipa1l2 Rasgrp2 Cyth3 Cat Cmya5 Dmd Akap7 Tgfb2 Pde4d Ecm1 Igf1r Tlr4 Itgb3

GO:0002376	immune system process	3.03E-06	0.0213073	1.67E+00	Klf2 Hc Slc46a2 Il7r Tnfr3 Ptprz1 Selp Cd27 Lat2 Ccr2 Epas1 Tgfb2 Procr Tspan2 Foxj1 Sptb Il18r1 Bcl6 Gcnt1 Gpc3 Cxcl3 Cd24a Tcf7 Pde4b Ms4a1 S1pr1 Sla2 Tlx1 Cxcr4 Ifnlr1 Txk P2ry14 Pigr Tdgr1 Slc8a3 Ctse Ptprz1 Cd300lf Cx3cr1 Pde4d Fasl Myb Samhd1 Tspan2 Gab3 Efna2 Tnfr3 Cx3cr1 Rora Pglyrp1 Gcnt1 Ifitm1 Il18r1 Foxp1 Pde4b Tcf7 Eml1 Spib Klf4 Serpinb9 Eml1 Ifitm1 Aicda Eomes Cfhlfnar1 Ifnar1 Pde4b Tgfb2 Pde4d Cblb Runx2 Igf1r Tlr4
GO:0004896 GO:0004907	cytokine receptor activity	3.25E-06	0.0213073	1.67E+00	Il7r Ccr2 Il1r2 Il18r1 Il18rap Cxcr4 Ifnlr1 Cxcr6 Il12rb2 Cx3cr1 Cxcr6 Ifnar1

GO:0009653	anatomical structure morphogenesis	5.97E-06	0.0339258	1.47E+00	Cacna1s Rapgef3 Tgfb3 Klf2 Serpinb5 Il7r Ptprz1 Selp Ptprb Ccr2 Epas1 Nyap2 mod1 Tgfb2 Ecm1 Foxj1 Sfrp4 Spaca1 Matn2 Hgf Col4a2 Bcl6 Ramp1 Vcl Gcnt1 Map2 Gpc3 Sema4f Tcf7 Ablim1 Nr4a2 S1pr1 Itgb3 Tlx1 Cxcr4 Murc Dcc Tdgf1 Atp2b2 Spaca1 Sall3 Gcm1 Ptprz1 Nr2e1 Cryaa Tead1 Wwox Fasl Cap2 Efna2 Fgfr1 Tbx6 Dusp5 Rora Gcnt1 Ifitm1 Hmx2 Foxp1 Adora2a Tcf7 Ctnnd2 Tmem106b Smo Klf4 Slitrk4 Ermn Flrt3 Ifitm1 Gcm1 Sema6c Thsd7a Ntn1 Dmd Eomes Runx2 Igf1r Crispld1
GO:0005891	voltage-gated calcium channel complex	6.66E-06	0.0339258	1.47E+00	Cacna1s Pde4b Catsper3 Cacna1h Pde4d Cacnb2 Catsperd

GO:0051239	regulation of multicellular organismal process	8.09E-06	0.0371204	1.43E+00	Dusp10 Rapgef3 Tgfb3 Cysltr2 Cd59a Klf2 Hc Slc46a2 Il7r Ptpz1 Selp Lama4 Cd27 Ccr2 Epas1 Aspa Lag3 Sytl2 Proc Xdh Tgfb2 Il20ra Procr Ecm1 Foxj1 Sfrp4 Prnp Hgf Col4a2 Il18r1 Tg Bcl6 Gpc3 Sema4f Cd24a Gcnt2 Pde4b Nr4a2 Ntrk3 S1pr1 Itgb3 Cxcr4 Hes5 Txk Dcc Tdgf1 Il12rb2 Grm1 Ptpz1 Scn3b Nr2e1 Mfap4 Ntrk3 Cx3cr1 Pde4d Fasl Myb Tnfaip3 Fgfr1 Sgip1 Tbx6 C1qtnf3 Rora Pglyrp1 Tnnt3 Adra1b P2rx4 Foxp1 Pde4b Adora2a Sipal1l1 Smo Klf4 Cmya5 Ntn1 Dmd Btg1 Eomes Kcnma1 Ifnar1 Cblb Runx2 Ccnd1 Tlr4
GO:0007166	cell surface receptor signaling pathway	8.93E-06	0.0372139	1.43E+00	Tgfb3 Il7r Wisp1 Cd27 Lat2 Ccr2 Lag3 Clnk Tgfb2 Il20ra Sfrp4 Il1r2 Hgf Il18r1 Fzd10 Cd24a Tcf7 Gcnt2 Pde4b Ntrk3 Il18rap Itgb3 Cxcr4 Ifnlr1 Hes5 Abca1 Plaur Txk P2ry14 Pigr Dcc Cxcr6 Tdgf1 Adamts14 Il12rb2 Grm1 Gria4 Adam34 Ntrk3 Cx3cr1 Wwox Fasl Kremen1 Efna2 Cx3cr1 Ntrk3 Fgfr1 Vipr1 Tle2 P2rx4 Pde4b Adora2a Tcf7 Sipal1l1 Smo Wisp1 Eya2 Gpr114 Cxcr6 Akap7 Ifnar1 Cblb Runx2 Igf1r Ccnd1 Tlr4

GO:0048583	regulation of response to stimulus	1.02E-05	0.0389147	1.41E+00	Dusp10 Rapgef3 Tgfb3 Cysltr2 Cd59a Hc Phlpp1 Il7r Tnip3 Abcb9 Selp Cd27 Lat2 Ccr2 Sesn3 Lag3 Proc Xdh Tgfb2 Il20ra Grb7 Foxj1 Sfrp4 Prnp Hgf Il18r1 Bcl6 Ramp1 Fzd10 Gpc3 Cd24a Gcnt2 Pde4b Fam110c Ntrk3 S1pr1 Itgb3 Sla2 Nt5e Cxcr4 Ifnlr1 Hes5 Abca1 Txk Ramp3 Pigr Dcc Tdgf1 Agpat9 Rasa3 Bicc1 Sall3 Grm1 Dusp7 Cx3cr1 Sell Pde4d Wwox Fasl Myb Samhd1 Tnfaip3 Fgfr1 Zfyve28 C1qtnf3 Dusp5 Padi2 Rora Pglyrp1 Tle2 Adra1b Ramp3 P2rx4 Adora2a Sipa1l1 Klf4 Sipa1l2 Wwc1 Rasgrp2 Cyth3 Cat Cmya5 Dmd Akap7 Cfh Pde4d Cblb Runx2 Ecm1 Igf1r Ccnd1 Kctd12 Tlr4
GO:0098552	side of membrane	1.38E-05	0.0486405	1.31E+00	Cd59a Il7r Selp Cd27 Ccr2 Lag3 Cd24a Klrc1 Ms4a1 S1pr1 Itgb3 Cxcr4 Abca1 Rasa3 Il12rb2 Klra4 Klra1 Klrc1 Sell Fasl Klrd1 Cyth3 Klrc3 Kcnma1 Cd33 Tlr4

SUPPLEMENTARY FIGURES LEGENDS

Figure E1. Effects of COX inhibitors on *Il23r* expression. (A) Expression of *Il23r* in differentiated Th17 cells treated with IL-23 (10 ng/ml) and/or PGE₂ (100 nM) in the absence or presence of a COX-1 inhibitor, SC-560 (100 µM) or a COX-2 inhibitor, SC-236 (100 µM) or both for 3 days. (B) Th17 cells were cultured with vehicle, IL-23 (10 ng/ml) and PGE₂ (100 nM) or IL-23 (10 ng/ml), EP2 agonist (100 nM) and EP4 agonist (100 nM) in the absence or presence of indomethacin (100 µM) for 3 days, and then harvested to analyze for *Il23r* expression by qRT-PCR.

Figure E2. db-cAMP activates JAK2 and STAT3 in Th17 cells. (A) Time-course of STAT3 Y705 phosphorylation by vehicle, db-cAMP, IL-23 or db-cAMP and IL-23 in Th17 cells. Th17 cells were treated with IL-23 (10 ng/ml) for 3 days to induce IL-23R expression. The cells were then stimulated with either 100 µM db-cAMP or 100 ng/ml IL-23 or in combination for indicated times. Phosphorylation of STAT3 at Y705 residues and S727 residues under each condition was examined at indicated times by Western blot analysis using total cell lysates and antibodies to each phosphorylation site

of STAT3, total STAT3 and α -Tubulin. Representative images are shown (n=2). (B)

Involvement of Src family kinase in cAMP-induced JAK2 Y1007/Y1008

phosphorylation. Th17 cells were stimulated with 100 μ M db-cAMP for 10 min in the

presence of a Src inhibitor, Src Kinase Inhibitor I (10 μ M), and subjected to Western

blot analysis. Data was from a single experiment.

Figure E3. Heat-map and gene ontology analysis of genes in each cluster. (A) Gene

expression profiles in Th17 cells stimulated with db-cAMP and/or IL-23 followed by

microarray analysis. Heat-map analysis of expression of genes 2-folds up- or

down-regulated upon each stimulus compared to the vehicle control (One-way ANOVA

$p < 0.05$, n=3). (B) Gene ontology analysis of each clusters by GeneSpring.

Figure E4. Involvement of PGE₂ signaling in psoriasis-like model. (A) Gene

expression of PGE₂ synthases in ear skin from naïve WT mice or psoriasis-like skin

lesions from mice administrated with IL-23 by intradermal injection in the dorsum (n=5

each). Gene expression was retrieved from a public dataset GSE13335. (B and C)

Genetic loss of *Ptger2* (EP2 KO) or pharmacological EP4 antagonism alone does not cause alteration in the ear. Psoriasis-like model in WT and EP2 KO mice were established as described in Figure 5A. Ear swelling was measured every 2 days (B) (n=14, 10, 8, and 10 in vehicle-treated mice, EP2 KO mice, AS1954813-treated WT mice, AS1954813-treated EP2 KO mice, respectively) and ear skins were subjected to FACS analysis on day 4 (C). (D and E) Suppression of IL-17A⁺ and IL-17A⁺IFN- γ ⁺ CD4⁺ T cell accumulation by EP2 KO and EP4 antagonist. WT and EP2 KO mice were administered either vehicle or AS1954813, and subcutaneously injected IL-23. CD4⁺ T cells were purified from the ear of each group on day 4 and examined by FACS for IL-17A and IFN- γ . Representative data from 4 independent experiments are shown. All bars indicate mean \pm SEM. *, p<0.05, **, p<0.01, ***, p<0.001.

Figure E5. FACS analysis of cell populations of EP2^{fl/fl}EP4^{fl/fl}Lck-Cre⁺ mice or WT Lck-Cre⁺ mice. (A) Cell population of EP2^{fl/fl}EP4^{fl/fl}Lck-Cre⁺ mice and control WT Lck-Cre⁺ mice. The numbers of B cell, T cell, CD4 T cell, CD8 T cell, Th1 cell, Th17 cell and Treg cell isolated from thymus, spleen, lymph node, and peripheral blood were

analyzed by FACS. (n=3-4)

Figure E6. Involvement of PGE₂ signaling in IMQ-induced psoriasis-like model and the effect of COX inhibitors on IL-23-induced psoriasis model. (A) Control WT mice and *Ptger2*-deficient (EP2 KO) mice were subjected to imiquimod (IMQ)-induced psoriasis model and administered either vehicle or AS1954813, 100 mg/kg, as described in Figure 5A. Ear swelling was measured every 2 days (A) (n=14, 10, 8, and 10 in vehicle-treated mice, EP2 KO mice, AS1954813-treated WT mice, AS1954813-treated EP2 KO mice, respectively). (B-D) Female WT mice were subjected to IL23-induced psoriasis model and administered either vehicle, SC-236 (10mg/kg) or indomethacin (4 mg/kg). Ear swelling was measured every 2 days (B) (n=4, respectively) and mice were sacrificed and subjected to FACS analysis at day 4 (C and D).

Figure E7. IL-1 β -IL1 receptor signaling was not involved in *Il23r* expression by Th17 cells. Expression of *Il23r* gene in differentiated Th17 cells stimulated with db-cAMP, IL-23 or db-cAMP and IL-23 in combination in the absence or presence of

various concentrations of neutralization antibody for IL-1 β for 3 days was analyzed by

qRT-PCR (n=3). All bars indicate mean \pm SEM.

Table E 1. **List of genes in Cluster 1U.**

Table E 2. **List of genes in Cluster 2U.**

Table E 3. **List of genes in Cluster 3U.**

Table E 4. **List of genes in Cluster 4U.**

Table E 5. **List of genes in Cluster 1D.**

Table E 6. **List of genes in Cluster 2D.**

Table E 7. **List of genes in Cluster 3D.**

81

82 Table E 8. **List of genes in Cluster 4D.**

83

84 Table E 9. **List of gene ontology from Cluster 1U.**

85

86 Table E 10. **List of gene ontology from Cluster 2U.**

87

88 Table E 11. **List of gene ontology from Cluster 3U.**




Figure E1




Figure E2




Figure E3




Figure E4

A





Figure E5




Figure E6




Figure E7